

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
27 February 2003 (27.02.2003)

PCT

(10) International Publication Number  
**WO 03/016306 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 403/12**,  
401/12, 407/12, 409/12, 413/12, 417/12, 403/14, 233/96,  
A61K 31/505, C07D 239/46, 239/48, C07C 211/00

(21) International Application Number: PCT/EP02/08953

(22) International Filing Date: 9 August 2002 (09.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
01203090.4 13 August 2001 (13.08.2001) EP  
02077748.8 10 June 2002 (10.06.2002) EP

(71) Applicant (for all designated States except US):  
**JANSSEN PHARMACEUTICA N.V.** [BE/BE]; Turn-  
houtseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GUILLEMONT, Jérôme, Emile, Georges** [FR/FR]; Janssen-Cilag S.A.,  
1, rue Camille Desmoulins, TSA 91003, F-92787  
Issy-les-Moulineaux (FR). **PALANDJIAN, Patrice**  
[FR/FR]; Janssen-Cilag S.A., 1, rue Camille Desmoulins,  
TSA 91003, F-92787 Issy-les-Moulineaux (FR). **DE JONGE, Marc, René** [NL/NL]; Janssen Pharmaceutica  
N.V., Turnhoutseweg 30, B-2340 Beerse (BE). **KOY-  
MANS, Lucien, Maria, Henricus** [NL/BE]; Janssen  
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse

(BE). **VINKERS, Hendrik, Maarten** [NL/BE]; Janssen  
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse  
(BE). **DAEYAERT, Frederik, Frans, Desiré** [BE/BE];  
Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340  
Beerse (BE). **HEERES, Jan** [NL/BE]; Janssen Pharma-  
ceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).  
**VAN AKEN, Koen, Jeanne, Alfons** [BE/BE]; Bruyn-  
ingstraat 3, B-8500 Kortrijk (BE). **LEWI, Paulus, Joannes**  
[BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg  
30, B-2340 Beerse (BE). **JANSSEN, Paul, Adriaan, Jan**  
[BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30,  
B-2340 Beerse (BE).


(74) Agents: **QUAGHEBEUR, Luc** et al.; Janssen Pharmaceu-  
tica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: HIV INHIBITING PYRIMIDINES DERIVATIVES



(57) Abstract: This invention concerns HIV replication inhibitors of formula (I) the *N*-oxides, the pharmaceutically acceptable addition salts, the quaternary amines and the stereochemically isomeric forms thereof, wherein the ring containing -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- and -b<sup>1</sup>=b<sup>2</sup>-b<sup>3</sup>=b<sup>4</sup>- represents phenyl, pyridyl, pyrimidinyl, pirazinyl, pyridazinyl; n is 0 to 5; m is 1 to 4; R<sup>1</sup> is hydrogen; aryl; formyl; C<sub>1-6</sub>alkylcarbonyl; C<sup>1-6</sup>alkyl; C<sub>1-6</sub>alkyloxycarbonyl, substituted C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl; R<sup>2</sup> is hydroxy, halo, optionally substituted C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl, optionally substituted C<sub>2-6</sub>alkenyl, optionally substituted C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl,

carboxyl, cyano, nitro, amino, mono- or di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sup>2</sup>?, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup> or a 5-membered heterocycle; X<sub>1</sub> is -NR<sup>5</sup>-, -NH-NH-, -N=N-, -O-, -C(=O)-, C<sub>1-4</sub>alkanediyl, -CHOH-, -S-, -S(=O)<sub>p</sub>-, -X<sub>2</sub>-C<sub>1-4</sub>alkanediyl- or -C<sub>1-4</sub>alkanediyl-X<sub>2</sub>-; R<sup>3</sup> is NHR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>; -C(=O)-NHR<sup>13</sup>; -C(=O)-NR<sup>13</sup>R<sup>14</sup>; -C(=O)-R<sup>15</sup>; -CH=N-NH-C(=O)-R<sup>16</sup>; substituted C<sub>1-6</sub>alkyl; optionally substituted C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl; substituted C<sub>2-6</sub>alkenyl; substituted C<sub>2-6</sub>alkynyl; C<sub>1-6</sub>alkyl substituted with hydroxy and a second substituent; -C(=N-O-R<sup>8</sup>)-C<sub>1-4</sub>alkyl; R<sup>7</sup>; or -X<sub>3</sub>-R<sup>7</sup>; R<sup>4</sup> is halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylcarbonyl, formyl, amino, mono- or di(C<sub>1-4</sub>alkyl)amino; their use as a medicine, their processes for preparation and pharmaceutical compositions comprising them.

WO 03/016306 A1

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB,

GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

**Published:**

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

## HIV INHIBITING PYRIMIDINES DERIVATIVES

The present invention is concerned with pyrimidine derivatives having HIV (Human Immunodeficiency Virus) replication inhibiting properties. The invention further relates to methods for their preparation and pharmaceutical compositions comprising them. The invention also relates to the use of said compounds for the manufacture of a medicament for the prevention or the treatment of HIV infection.

Compounds structurally related to the present compounds are disclosed in the prior art.

WO 99/50250 and WO 00/27825 disclose substituted aminopyrimidines having HIV replication inhibiting properties.

WO 97/19065 discloses substituted 2-anilinopyrimidines useful as protein kinase inhibitors.

WO 00/62778 concerns cyclic protein tyrosine kinase inhibitors.

WO 98/41512 describes substituted 2-anilinopyrimidines useful as protein kinase inhibitors.


US 5,691,364 describes benzamidine derivatives and their use as anti-coagulants.

WO 00/78731 describes 5-cyano-2-aminopyrimidine derivatives as KDR kinase or FGFR kinase inhibitors useful in the prophylaxis and treatment of diseases associated with angiogenesis.

The compounds of the invention differ from the prior art compounds in structure, pharmacological activity and/or pharmacological potency.

Unexpectedly, it has been found that the compounds of the invention have an improved ability to inhibit the replication of Human Immunodeficiency Virus (HIV), in particular they have an improved ability to inhibit the replication of mutant strains, i.e. strains which have become resistant to art-known drug(s) (drug or multidrug resistant HIV strains).

The present invention concerns a compound of formula



-2-

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein

$-a^1=a^2-a^3=a^4-$  represents a bivalent radical of formula

- 5       $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$       (a-1);  
        $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$       (a-2);  
        $-\text{N}=\text{CH}-\text{N}=\text{CH}-$       (a-3);  
        $-\text{N}=\text{CH}-\text{CH}=\text{N}-$       (a-4);  
        $-\text{N}=\text{N}-\text{CH}=\text{CH}-$       (a-5);

$-b^1=b^2-b^3=b^4-$  represents a bivalent radical of formula

- 10       $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$       (b-1);  
        $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$       (b-2);  
        $-\text{N}=\text{CH}-\text{N}=\text{CH}-$       (b-3);  
        $-\text{N}=\text{CH}-\text{CH}=\text{N}-$       (b-4);  
        $-\text{N}=\text{N}-\text{CH}=\text{CH}-$       (b-5);

15      *n* is 0, 1, 2, 3 or 4; and in case  $-a^1=a^2-a^3=a^4-$  is (a-1), then *n* may also be 5;

*m* is 1, 2, 3 and in case  $-b^1=b^2-b^3=b^4-$  is (b-1), then *m* may also be 4;

*R*<sup>1</sup> is hydrogen; aryl; formyl; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxycarbonyl;

C<sub>1-6</sub>alkyl substituted with formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl,

C<sub>1-6</sub>alkylcarbonyloxy; C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkylcarbonyl substituted with

20      C<sub>1-6</sub>alkyloxycarbonyl;


each *R*<sup>2</sup> independently is hydroxy, halo, C<sub>1-6</sub>alkyl optionally substituted with cyano or

$-\text{C}(=\text{O})\text{R}^6$ , C<sub>3-7</sub>cycloalkyl, C<sub>2-6</sub>alkenyl optionally substituted with one or more

halogen atoms or cyano, C<sub>2-6</sub>alkynyl optionally substituted with one or more halogen

atoms or cyano, C<sub>1-6</sub>alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or

25      di(C<sub>1-6</sub>alkyl)amino, polyhalomethyl, polyhalomethylthio,  $-\text{S}(=\text{O})_p\text{R}^6$ ,  $-\text{NH}-\text{S}(=\text{O})_p\text{R}^6$ ,  
 $-\text{C}(=\text{O})\text{R}^6$ ,  $-\text{NHC}(=\text{O})\text{H}$ ,  $-\text{C}(=\text{O})\text{NHNH}_2$ ,  $-\text{NHC}(=\text{O})\text{R}^6$ ,  $-\text{C}(=\text{NH})\text{R}^6$  or a radical of  
 formula



wherein each A<sub>1</sub> independently is N, CH or CR<sup>6</sup>; and

30      A<sub>2</sub> is NH, O, S or NR<sup>6</sup>;

X<sub>1</sub> is  $-\text{NR}^5-$ ,  $-\text{NH}-\text{NH}-$ ,  $-\text{N}=\text{N}-$ ,  $-\text{O}-$ ,  $-\text{C}(=\text{O})-$ , C<sub>1-4</sub>alkanediyl,  $-\text{CHOH}-$ ,  $-\text{S}-$ ,  $-\text{S}(=\text{O})_p-$ ,  
 $-\text{X}_2-\text{C}_{1-4}\text{alkanediyl}-$  or  $-\text{C}_{1-4}\text{alkanediyl}-\text{X}_2-$ ;

X<sub>2</sub> is  $-\text{NR}^5-$ ,  $-\text{NH}-\text{NH}-$ ,  $-\text{N}=\text{N}-$ ,  $-\text{O}-$ ,  $-\text{C}(=\text{O})-$ ,  $-\text{CHOH}-$ ,  $-\text{S}-$ ,  $-\text{S}(=\text{O})_p-$ ;

*R*<sup>3</sup> is  $\text{NHR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $-\text{C}(=\text{O})-\text{NHR}^{13}$ ,  $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$ ,  $-\text{C}(=\text{O})-\text{R}^{15}$ ,  $-\text{CH}=\text{N}-\text{NH}-$

35       $\text{C}(=\text{O})-\text{R}^{16}$ ; C<sub>1-6</sub>alkyl substituted with one or more substituents each

independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or

-3-

- $R^7$ ;  $C_{1-6}$ alkyl substituted with one or more substituents each independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  $C_{1-4}$ alkanediyl;  $C_{1-6}$ alkyl substituted with hydroxy and a second substituent selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl optionally substituted with one or more substituents each independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{2-6}$ alkenyl substituted with one or more substituents each independently selected from halo, cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{2-6}$ alkynyl substituted with one or more substituents each independently selected from halo, cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $-C(=N-O-R^8)-C_{1-4}$ alkyl;  $R^7$  or  $-X_3-R^7$ ;
- $X_3$  is  $-NR^5-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-C(=O)-$ ,  $-S-$ ,  $-S(=O)_p-$ ,  $-X_2-C_{1-4}$ alkanediyl-,  $-C_{1-4}$ alkanediyl- $X_{2a}-$ ,  $-C_{1-4}$ alkanediyl- $X_{2b}-C_{1-4}$ alkanediyl-,  $-C(=N-OR^8)-C_{1-4}$ alkanediyl-;
- with  $X_{2a}$  being  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-C(=O)-$ ,  $-S-$ ,  $-S(=O)_p-$ ; and  
 with  $X_{2b}$  being  $-NH-NH-$ ,  $-N=N-$ ,  $-C(=O)-$ ,  $-S-$ ,  $-S(=O)_p-$ ;
- $R^4$  is halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy, cyano, nitro, polyhalo $C_{1-6}$ alkyl, polyhalo $C_{1-6}$ alkyloxy, aminocarbonyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl, formyl, amino, mono- or di( $C_{1-4}$ alkyl)amino or  $R^7$ ;
- $R^5$  is hydrogen; aryl; formyl;  $C_{1-6}$ alkylcarbonyl;  $C_{1-6}$ alkyl;  $C_{1-6}$ alkyloxycarbonyl;  $C_{1-6}$ alkyl substituted with formyl,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyloxycarbonyl or  $C_{1-6}$ alkylcarbonyloxy;  $C_{1-6}$ alkyloxy $C_{1-6}$ alkylcarbonyl substituted with  $C_{1-6}$ alkyloxycarbonyl;
- $R^6$  is  $C_{1-4}$ alkyl, amino, mono- or di( $C_{1-4}$ alkyl)amino or polyhalo $C_{1-4}$ alkyl;
- $R^7$  is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl, mono or di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl, formyl,  $C_{1-6}$ alkylcarbonyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylthio, cyano, nitro, polyhalo $C_{1-6}$ alkyl, polyhalo $C_{1-6}$ alkyloxy, aminocarbonyl,  $-CH(=N-O-R^8)$ ,  $R^{7a}$ ,  $-X_3-R^{7a}$  or  $R^{7a}-C_{1-4}$ alkyl;
- $R^{7a}$  is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems

-4-

may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, aminoC<sub>1-6</sub>alkyl, mono or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylthio, cyano, nitro,

5 polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl, -CH(=N-O-R<sup>8</sup>);

R<sup>8</sup> is hydrogen, C<sub>1-4</sub>alkyl, aryl or arylC<sub>1-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently are hydrogen; hydroxy; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxy;

C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; amino; mono- or di(C<sub>1-6</sub>alkyl)amino;

mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl; -CH(=NR<sup>11</sup>) or R<sup>7</sup>, wherein each of the






10 aforementioned C<sub>1-6</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy,

C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, mono- or di(C<sub>1-4</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy,

polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H,

15 -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup>, R<sup>7</sup>; or

R<sup>9</sup> and R<sup>10</sup> may be taken together to form a bivalent or trivalent radical of formula



R<sup>11</sup> is cyano; C<sub>1-4</sub>alkyl optionally substituted with C<sub>1-4</sub>alkyloxy, cyano, amino, mono- or

25 di(C<sub>1-4</sub>alkyl)amino or aminocarbonyl; C<sub>1-4</sub>alkylcarbonyl; C<sub>1-4</sub>alkyloxycarbonyl; aminocarbonyl; mono- or di(C<sub>1-4</sub>alkyl)aminocarbonyl;

R<sup>12</sup> is hydrogen or C<sub>1-4</sub>alkyl;

R<sup>13</sup> and R<sup>14</sup> each independently are C<sub>1-6</sub>alkyl optionally substituted with cyano or aminocarbonyl, C<sub>2-6</sub>alkenyl optionally substituted with cyano or aminocarbonyl,

30 C<sub>2-6</sub>alkynyl optionally substituted with cyano or aminocarbonyl;

R<sup>15</sup> is C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;

R<sup>16</sup> is C<sub>1-6</sub>alkyl optionally substituted with cyano or aminocarbonyl, or R<sup>7</sup>;

p is 1 or 2;

aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each

35 independently selected from halo, hydroxy, mercapto, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, aminoC<sub>1-6</sub>alkyl, mono or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl,

-5-

C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylthio, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl, R<sup>7</sup> or -X<sub>3</sub>-R<sup>7</sup>.

As used hereinbefore or hereinafter C<sub>1-4</sub>alkyl as a group or part of a group defines  
 5 straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl; C<sub>1-6</sub>alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the group defined for C<sub>1-4</sub>alkyl and pentyl, hexyl, 2-methylbutyl and the like; C<sub>2-6</sub>alkyl as a group or part of a group defines straight or  
 10 branched chain saturated hydrocarbon radicals having from 2 to 6 carbon atoms such as ethyl, propyl, 1-methylethyl, butyl, pentyl, hexyl, 2-methylbutyl and the like; C<sub>1-4</sub>alkanediyl defines straight or branched chain saturated bivalent hydrocarbon radicals having from 1 to 4 carbon atoms such as methylene, 1,2-ethanediyl or 1,2-ethylidene, 1,3-propanediyl or 1,3-propylidene, 1,4-butanediyl or  
 15 1,4-butylidene and the like; C<sub>3-7</sub>cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; C<sub>2-6</sub>alkenyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms containing a double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl and the like; C<sub>2-6</sub>alkynyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms  
 20 containing a triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like; a monocyclic, bicyclic or tricyclic saturated carbocycle represents a ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms and said ring system containing only single bonds; a monocyclic, bicyclic or tricyclic partially saturated carbocycle represents a ring system consisting of 1, 2 or 3 rings, said  
 25 ring system being composed of only carbon atoms and comprising at least one double bond provided that the ring system is not an aromatic ring system; a monocyclic, bicyclic or tricyclic aromatic carbocycle represents an aromatic ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms; the term aromatic is well known to a person skilled in the art and designates cyclically  
 30 conjugated systems of 4n + 2 electrons, that is with 6, 10, 14 etc.  $\pi$ -electrons (rule of Hückel) ; a monocyclic, bicyclic or tricyclic saturated heterocycle represents a ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected from O, N or S, said ring system containing only single bonds; a monocyclic, bicyclic or tricyclic partially saturated heterocycle represents a ring system consisting of 1, 2 or  
 35 3 rings and comprising at least one heteroatom selected from O, N or S, and at least one double bond provided that the ring system is not an aromatic ring system; a monocyclic,

-6-

bicyclic or tricyclic aromatic heterocycle represents an aromatic ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected from O, N or S.

5 Particular examples of monocyclic, bicyclic or tricyclic saturated carbocycles are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[4,2,0]octanyl, cyclononanyl, cyclodecanyl, decahydronaphthalenyl, tetradecahydroanthracenyl and the like.

10 Particular examples of monocyclic, bicyclic or tricyclic partially saturated carbocycles are cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, bicyclo[4,2,0]octenyl, cyclononenyl, cyclodecenyl, octahydronaphthalenyl, 1,2,3,4-tetrahydronaphthalenyl, 1,2,3,4,4a,9,9a,10-octahydro-anthracenyl and the like.

15 Particular examples of monocyclic, bicyclic or tricyclic aromatic carbocycles are phenyl, naphthalenyl, anthracenyl.

20 Particular examples of monocyclic, bicyclic or tricyclic saturated heterocycles are tetrahydrofuranyl, pyrrolidinyl, dioxolanyl, imidazolidinyl, thiazolidinyl, tetrahydrothienyl, dihydrooxazolyl, isothiazolidinyl, isoxazolidinyl, oxadiazolidinyl, triazolidinyl, thiadiazolidinyl, pyrazolidinyl, piperidinyl, hexahydropyrimidinyl, hexahydropyrazinyl, dioxanyl, morpholinyl, dithianyl, thiomorpholinyl, piperazinyl, trithianyl, decahydroquinolinyl, octahydroindolyl and the like.

25 Particular examples of monocyclic, bicyclic or tricyclic partially saturated heterocycles are pyrrolinyl, imidazolinyl, pyrazolinyl, 2,3-dihydrobenzofuranyl, 1,3-benzodioxolyl, 2,3-dihydro-1,4-benzodioxinyl, indolinyl and the like.

30 Particular examples of monocyclic, bicyclic or tricyclic aromatic heterocycles are azetyl, oxetylidenyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, benzofuryl, isobenzofuryl, benzothienyl, isobenzothienyl, indolizinyl, indolyl, isoindolyl, benzoxazolyl, benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benzopyrazolyl, benzoxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl, 35 cinnolinyl, quinolizinyl, phthalazinyl, quinoxalinyl, quinazolinyl, naphthiridinyl, pteridinyl, benzopyranyl, pyrrolopyridyl, thienopyridyl, furopyridyl, isothiazolopyridyl, thiazolopyridyl, isoxazolopyridyl, oxazolopyridyl, pyrazolopyridyl, imidazopyridyl,



-7-

pyrrolopyrazinyl, thienopyrazinyl, furopyrazinyl, isothiazolopyrazinyl,  
thiazolopyrazinyl, isoxazolopyrazinyl, oxazolopyrazinyl, pyrazolopyrazinyl,  
imidazopyrazinyl, pyrrolopyrimidinyl, thienopyrimidinyl, furopyrimidinyl,  
isothiazolopyrimidinyl, thiazolopyrimidinyl, isoxazolopyrimidinyl, oxazolopyrimidinyl,  
5 pyrazolopyrimidinyl, imidazopyrimidinyl, pyrrolopyridazinyl, thienopyridazinyl,  
furopyridazinyl, isothiazolopyridazinyl, thiazolopyridazinyl, isoxazolopyridazinyl,  
oxazolopyridazinyl, pyrazolopyridazinyl, imidazopyridazinyl, oxadiazolopyridyl,  
thiadiazolopyridyl, triazolopyridyl, oxadiazolopyrazinyl, thiadiazolopyrazinyl,  
triazolopyrazinyl, oxadiazolopyrimidinyl, thiadiazolopyrimidinyl, triazolopyrimidinyl,  
10 oxadiazolopyridazinyl, thiadiazolopyridazinyl, triazolopyridazinyl, imidazo-oxazolyl,  
imidazothiazolyl, imidazoimidazolyl, isoxazolotriazinyl, isothiazolotriazinyl,  
pyrazolotriazinyl, oxazolotriazinyl, thiazolotriazinyl, imidazotriazinyl,  
oxadiazolotriazinyl, thiadiazolotriazinyl, triazolotriazinyl, carbazolyl, acridinyl,  
phenazinyl, phenothiazinyl, phenoxazinyl and the like.

15

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom.

20

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhalomethyl as a group or part of a group is defined as mono- or polyhalosubstituted methyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl; polyhaloC<sub>1-4</sub>alkyl or polyhaloC<sub>1-6</sub>alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C<sub>1-4</sub>alkyl or  
25 C<sub>1-6</sub>alkyl, for example, the groups defined in halomethyl, 1,1-difluoro-ethyl and the like. In case more than one halogen atoms are attached to an alkyl group within the definition of polyhalomethyl, polyhaloC<sub>1-4</sub>alkyl or polyhaloC<sub>1-6</sub>alkyl, they may be the same or different.

30

The term heterocycle in the definition of R<sup>7</sup> or R<sup>7a</sup> is meant to include all the possible isomeric forms of the heterocycles, for instance, pyrrolyl comprises 1*H*-pyrrolyl and 2*H*-pyrrolyl.

35

The carbocycle or heterocycle in the definition of R<sup>7</sup> or R<sup>7a</sup> may be attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate, if not otherwise specified. Thus, for example, when the heterocycle is

-8-

imidazolyl, it may be 1-imidazolyl, 2-imidazolyl, 4-imidazolyl and the like, or when the carbocycle is naphthalenyl, it may be 1-naphthalenyl, 2-naphthalenyl and the like.

- 5 When any variable (eg.  $R^7$ ,  $X_2$ ) occurs more than one time in any constituent, each definition is independent.

Lines drawn from substituents into ring systems indicate that the bond may be attached to any of the suitable ring atoms.

- 10 For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether  
15 pharmaceutically acceptable or not are included within the ambit of the present invention.

- The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. The latter can conveniently be obtained by  
20 treating the base form with such appropriate acids as inorganic acids, for example, hydrohalic acids, e.g. hydrochloric, hydrobromic and the like; sulfuric acid; nitric acid; phosphoric acid and the like; or organic acids, for example, acetic, propanoic, hydroxy-acetic, 2-hydroxypropanoic, 2-oxopropanoic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic,  
25 ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted by treatment with alkali into the free base form.

- The compounds of formula (I) containing acidic protons may be converted into their  
30 therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. primary, secondary and tertiary aliphatic and aromatic amines such as methylamine,  
35 ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine,

quinuclidine, pyridine, quinoline and isoquinoline, the benzathine, *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.

The term addition salt also comprises the hydrates and solvent addition forms which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several tertiary nitrogen atoms are oxidized to the so-called *N*-oxide.

It will be appreciated that some of the compounds of formula (I) and their *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms which the compounds of formula (I), and their *N*-oxides, addition salts, quaternary amines or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I) and their *N*-oxides, salts, solvates or quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other

-10-

isomers. Thus, when a compound of formula (I) is for instance specified as (E), this means that the compound is substantially free of the (Z) isomer.

In particular, stereogenic centers may have the R- or S-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the *cis*- or *trans*-

5 configuration. Compounds encompassing double bonds can have an E (entgegen) or Z (zusammen) -stereochemistry at said double bond. The terms *cis*, *trans*, R, S, E and Z are well known to a person skilled in the art.

Stereochemically isomeric forms of the compounds of formula (I) are obviously intended to be embraced within the scope of this invention.

10 For some of the compounds of formula (I), their prodrugs, *N*-oxides, salts, solvates, quaternary amines or metal complexes and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined. In these cases the stereoisomeric form which was first isolated is designated as "A" and the second as "B", without further reference to the actual stereochemical configuration.

15 However, said "A" and "B" stereoisomeric forms can be unambiguously characterized by for instance their optical rotation in case "A" and "B" have an enantiomeric relationship. A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction. In case "A" and "B" are stereoisomeric mixtures, they can be further separated whereby

20 the respective first fractions isolated are designated "A1" and "B1" and the second as "A2" and "B2", without further reference to the actual stereochemical configuration.

Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included

25 within the scope of the present invention.

Whenever used hereinafter, the term "compounds of formula (I)" is meant to also include their *N*-oxide forms, their salts, their quaternary amines and their stereochemically isomeric forms. Of special interest are those compounds of formula

30 (I) which are stereochemically pure.

Whenever used hereinbefore or hereinafter that substituents can be selected each independently out of a list of numerous definitions, such as for example for R<sup>9</sup> and R<sup>10</sup>, all possible combinations are intended which are chemically possible and which lead to

35 chemically stable molecules.


-11-

A particular group of compounds are those compounds of formula (I) wherein  $R^3$  is  $C_{1-6}$ alkyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{1-6}$ alkyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  $C_{1-4}$ alkanediyl;  $C_{1-6}$ alkyl substituted with hydroxy and a second substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{1-6}$ alkyloxy  $C_{1-6}$ alkyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{2-6}$ alkenyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{2-6}$ alkynyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $-C(=N-O-R^8)-C_{1-4}$ alkyl;  $R^7$  or  $-X_3-R^7$ ;  $R^4$  is halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy, cyano, nitro, polyhalo  $C_{1-6}$ alkyl, polyhalo  $C_{1-6}$ alkyloxy, aminocarbonyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl, formyl, amino, mono- or di( $C_{1-4}$ alkyl)amino;  $R^7$  is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto,  $C_{1-6}$ alkyl, hydroxy  $C_{1-6}$ alkyl, amino  $C_{1-6}$ alkyl, mono or di( $C_{1-6}$ alkyl)amino  $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylthio, cyano, nitro, polyhalo  $C_{1-6}$ alkyl, polyhalo  $C_{1-6}$ alkyloxy, aminocarbonyl,  $R^{7a}$ ,  $-X_3-R^{7a}$  or  $R^{7a}-C_{1-4}$ alkyl;  $R^{7a}$  is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto,  $C_{1-6}$ alkyl, hydroxy  $C_{1-6}$ alkyl, amino  $C_{1-6}$ alkyl, mono or di( $C_{1-6}$ alkyl)amino  $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylthio, cyano, nitro, polyhalo  $C_{1-6}$ alkyl, polyhalo  $C_{1-6}$ alkyloxy, aminocarbonyl;  $R^9$  and  $R^{10}$  each independently are hydrogen; hydroxy;  $C_{1-6}$ alkyl;  $C_{1-6}$ alkyloxy;  $C_{1-6}$ alkylcarbonyl;  $C_{1-6}$ alkyloxycarbonyl; amino; mono- or di( $C_{1-6}$ alkyl)amino; mono- or di( $C_{1-6}$ alkyl)aminocarbonyl or  $R^7$ , wherein each of the aforementioned  $C_{1-6}$ alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy,  $C_{1-6}$ alkyloxy, hydroxy  $C_{1-6}$ alkyloxy, carboxyl,  $C_{1-6}$ alkyloxycarbonyl, cyano, amino, imino, mono- or di( $C_{1-4}$ alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio,  $-S(=O)_pR^6$ ,  $-NH-S(=O)_pR^6$ ,  $-C(=O)R^6$ ,  $-NHC(=O)H$ ,  $-C(=O)NHNH_2$ ,  $-NHC(=O)R^6$ ,  $-C(=NH)R^6$ ,  $R^7$ .

-12-

An interesting group of compounds are those compounds of formula (I) wherein  $-a^1=a^2-a^3=a^4-$  represents a bivalent radical of formula  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$  (a-1).

- 5 Also an interesting group of compounds are those compounds of formula (I) having the formula




the *N*-oxides, the pharmaceutically acceptable addition salts, the quaternary amines or the stereochemically isomeric forms thereof, wherein

- 10  $-a^1=a^2-a^3=a^4-$ ,  $-b^1=b^2-b^3=b^4-$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $m$  and  $X_1$  are as defined hereinabove;  $n'$  is 0, 1, 2 or 3 and in case  $-a^1=a^2-a^3=a^4-$  is (a-1), then  $n'$  may also be 4;  $R^{2'}$  is halo,  $C_{1-6}$ alkyl, trihalomethyl, cyano, aminocarbonyl,  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl; provided that  $R^{2'}$  is placed at the para position in respect of the  $NR^1$  moiety.

15

Another interesting group of compounds are those compounds of formula (I) having the formula




the *N*-oxides, the pharmaceutically acceptable addition salts, the quaternary amines or the stereochemically isomeric forms thereof, wherein

- 20  $-b^1=b^2-b^3=b^4-$ ,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $m$  and  $X_1$  are as defined hereinabove;  $n'$  is 0, 1, 2, 3 or 4;  $R^{2'}$  is halo,  $C_{1-6}$ alkyl, trihalomethyl, cyano, aminocarbonyl,  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl.

25

Yet a further interesting group of compounds are those compounds of formula (I) having the formula



-13-

the *N*-oxides, the pharmaceutically acceptable addition salts, the quaternary amines or the stereochemically isomeric forms thereof, wherein

$R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $X_1$  are as defined hereinabove;


$n'$  is 0, 1, 2, 3 or 4;

- 5  $R^{2'}$  is halo,  $C_{1-6}$ alkyl, trihalomethyl, cyano, aminocarbonyl,  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl.

Also particular compounds are those compounds of formula (I), (I'), (I'') or (I''')

wherein one or wherever possible more of the following conditions apply :

- 10 a)  $m$  is 1, 2 or 3, in particular 2 or 3, more in particular 2, even more in particular  $m$  is 2 and said two  $R^4$  substituents are placed in position 2 and 6 (ortho position) in respect of the  $X_1$  moiety;
- b)  $m$  is 1, 2 or 3 and  $R^3$  is placed in position 4 (para position) in respect of the  $X_1$  moiety;
- 15 c)  $X_1$  is  $-NR^5-$ ,  $-NH-NH-$ ,  $-N=N-$ ,  $-O-$ ,  $-C(=O)-$ ,  $C_{1-4}$ alkanediyl,  $-CHOH-$ ,  $-S(=O)_p-$ ,  $-X_2-C_{1-4}$ alkanediyl- or  $-C_{1-4}$ alkanediyl- $X_2-$ ;
- d) where applicable  $n'$  is 0;
- e) where applicable  $n$  is 1 and said  $R^2$  substituent is placed in position 4 (para position) in respect of the  $NR^1$ -linker;
- 20 f)  $R^2$  is hydroxy, halo,  $C_{1-6}$ alkyl optionally substituted with cyano or  $-C(=O)R^6$ ,  $C_{3-7}$ cycloalkyl,  $C_{2-6}$ alkenyl optionally substituted with one or more halogen atoms or cyano,  $C_{2-6}$ alkynyl optionally substituted with one or more halogen atoms or cyano,  $C_{1-6}$ alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di( $C_{1-6}$ alkyl)amino, polyhalomethyl, polyhalomethylthio,  $-S(=O)_pR^6$ ,  $-NH-S(=O)_pR^6$ ,  $-NHC(=O)H$ ,  $-C(=O)NHNH_2$ ,  $-NHC(=O)R^6$ ,  $-C(=NH)R^6$  or a radical of formula
- 25



wherein each  $A_1$  independently is N, CH or  $CR^6$ ; and

$A_2$  is NH, O, S or  $NR^6$ ;

- g)  $R^{2'}$  is halo,  $C_{1-6}$ alkyl, trihalomethyl, cyano,  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl;
- 30 h)  $R^2$  is cyano, aminocarbonyl or  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl, in particular cyano;
- i)  $R^{2'}$  is cyano, aminocarbonyl or  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl, in particular cyano.
- 35

-14-

A preferred embodiment encompasses those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $NHR^{13}$ ,  $NR^{13}R^{14}$ ,  $-C(=O)-NHR^{13}$ ,  $-C(=O)-NR^{13}R^{14}$ ,  $-C(=O)-R^{15}$ ,  $-CH=N-NH-C(=O)-R^{16}$ ;  $C_{2-6}$ alkyl substituted with cyano or aminocarbonyl;  $C_{1-6}$ alkyl substituted with  $NR^9R^{10}$ ,  $-C(=O)-NR^{9a}R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{1-6}$ alkyl

- 5 substituted with two or more substituents each independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{1-6}$ alkyl substituted with one or more substituents each independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  $C_{1-4}$ alkanediyl;  $C_{1-6}$ alkyl substituted with hydroxy and a second
- 10 substituent selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl optionally substituted with one or more substituents each independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{2-6}$ alkenyl substituted with one or more substituents each independently selected from halo, cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{2-6}$ alkynyl
- 15 substituted with one or more substituents each independently selected from halo, cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $-C(=N-O-R^8)-C_{1-4}$ alkyl;  $R^7$  or  $-X_3-R^7$ ; with  $R^{9a}$  representing hydroxy;  $C_{1-6}$ alkyl;  $C_{1-6}$ alkyloxy;  $C_{1-6}$ alkylcarbonyl;  $C_{1-6}$ alkyloxycarbonyl; amino; mono- or di( $C_{1-6}$ alkyl)amino; mono- or di( $C_{1-6}$ alkyl)aminocarbonyl,  $-CH(=NR^{11})$  or  $R^7$ , wherein each of the aforementioned
- 20  $C_{1-6}$ alkyl groups in the definition of  $R^{9a}$  may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy,  $C_{1-6}$ alkyloxy, hydroxy $C_{1-6}$ alkyloxy, carboxyl,  $C_{1-6}$ alkyloxycarbonyl, cyano, amino, imino, mono- or di( $C_{1-4}$ alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio,  $-S(=O)_pR^6$ ,  $-NH-S(=O)_pR^6$ ,  $-C(=O)R^6$ ,  $-NHC(=O)H$ ,
- 25  $-C(=O)NHNH_2$ ,  $-NHC(=O)R^6$ ,  $-C(=NH)R^6$ ,  $R^7$ ;  $R^{9a}$  may also be taken together with  $R^{10}$  to form a bivalent or trivalent radical of formula (d-1), (d-2), (d-3), (d-4), (d-5), (d-6) or (d-7) as defined hereinabove.

- A further interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $NHR^{13}$ ,  $NR^{13}R^{14}$ ,  $-C(=O)-NHR^{13}$ ,  $-C(=O)-NR^{13}R^{14}$ ,  $-C(=O)-R^{15}$ ,  $-CH=N-NH-C(=O)-R^{16}$ ;  $C_{1-6}$ alkyl substituted with  $NR^9R^{10}$ ,  $-C(=O)-NR^{9a}R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{1-6}$ alkyl substituted with two or more substituents each
- 30 independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;  $C_{1-6}$ alkyl substituted with one or more substituents each independently selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  $C_{1-4}$ alkanediyl;  $C_{1-6}$ alkyl substituted
- 35 with hydroxy and a second substituent selected from cyano,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,



-15-

-C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl optionally substituted with one or more substituents each independently selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>,  
 -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>2-6</sub>alkenyl substituted with one or more substituents each independently selected from halo, cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>;  
 5 C<sub>2-6</sub>alkynyl substituted with one or more substituents each independently selected from halo, cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>;  
 -C(=N-O-R<sup>8</sup>)-C<sub>1-4</sub>alkyl; R<sup>7</sup> or -X<sub>3</sub>-R<sup>7</sup>; with R<sup>9a</sup> representing hydroxy; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl, -CH(=NR<sup>11</sup>) or R<sup>7</sup>, wherein  
 10 each of the aforementioned C<sub>1-6</sub>alkyl groups in the definition of R<sup>9a</sup> may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, mono- or di(C<sub>1-4</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>,  
 15 -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup>, R<sup>7</sup>; R<sup>9a</sup> may also be taken together with R<sup>10</sup> to form a bivalent or trivalent radical of formula (d-1), (d-2), (d-3), (d-4), (d-5), (d-6) or (d-7) as defined hereinabove.

Also an interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein R<sup>3</sup> is -CH=N-NH-C(=O)-R<sup>16</sup>; C<sub>1-6</sub>alkyl substituted with NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9a</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>1-6</sub>alkyl substituted with two or more substituents each independently selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>1-6</sub>alkyl substituted with one or more substituents each independently selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>  
 25 and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by C<sub>1-4</sub>alkanediyl; C<sub>1-6</sub>alkyl substituted with hydroxy and a second substituent selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl optionally substituted with one or more substituents each independently selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>2-6</sub>alkenyl substituted with  
 30 one or more substituents each independently selected from halo, cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; C<sub>2-6</sub>alkynyl substituted with one or more substituents each independently selected from halo, cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>; -C(=N-O-R<sup>8</sup>)-C<sub>1-4</sub>alkyl; R<sup>7</sup> or -X<sub>3</sub>-R<sup>7</sup>; with R<sup>9a</sup> as defined hereinabove.

35 Another interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein R<sup>3</sup> is NHR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>, -C(=O)-R<sup>15</sup>, C<sub>1-6</sub>alkyl substituted with one or

-16-

- more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{1-6}\text{alkyl}$  substituted with one or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by
- 5  $\text{C}_{1-4}\text{alkanediyl}$ ;  $\text{C}_{1-6}\text{alkyl}$  substituted with hydroxy and a second substituent selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{1-6}\text{alkyloxyC}_{1-6}\text{alkyl}$  optionally substituted with one or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{2-6}\text{alkenyl}$  substituted with one or more substituents each independently selected from halo, cyano,  $\text{NR}^9\text{R}^{10}$ ,
- 10  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{2-6}\text{alkynyl}$  substituted with one or more substituents each independently selected from halo, cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $-\text{C}(=\text{N}-\text{O}-\text{R}^8)-\text{C}_{1-4}\text{alkyl}$ ;  $\text{R}^7$  or  $-\text{X}_3-\text{R}^7$ .
- Also interesting are those compounds of formula (I), (I'), (I'') or (I''') wherein  $\text{R}^3$  is
- 15  $\text{C}_{1-6}\text{alkyl}$  substituted with  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^{9a}\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{1-6}\text{alkyl}$  substituted with two or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{1-6}\text{alkyl}$  substituted with one or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,
- 20  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  $\text{C}_{1-4}\text{alkanediyl}$ ;  $\text{C}_{1-6}\text{alkyl}$  substituted with hydroxy and a second substituent selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{1-6}\text{alkyloxyC}_{1-6}\text{alkyl}$  optionally substituted with one or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,
- 25  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{2-6}\text{alkenyl}$  substituted with one or more substituents each independently selected from halo, cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;  $\text{C}_{2-6}\text{alkynyl}$  substituted with one or more substituents each independently selected from halo, cyano,  $\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{NR}^9\text{R}^{10}$ ,  $-\text{C}(=\text{O})-\text{C}_{1-6}\text{alkyl}$  or  $\text{R}^7$ ;
- 30  $-\text{C}(=\text{N}-\text{O}-\text{R}^8)-\text{C}_{1-4}\text{alkyl}$ ;  $\text{R}^7$  or  $-\text{X}_3-\text{R}^7$ ; with  $\text{R}^{9a}$  as defined hereinabove.
- Also interesting are those compounds of formula (I), (I'), (I'') or (I''') wherein  $\text{R}^3$  is
- 35  $\text{C}_{1-6}\text{alkyl}$  substituted with one or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$  or  $\text{R}^7$ ;  $\text{C}_{2-6}\text{alkenyl}$  substituted with one or more substituents each independently selected from cyano,  $\text{NR}^9\text{R}^{10}$  or  $\text{R}^7$ ;  $\text{C}_{1-6}\text{alkyloxyC}_{1-6}\text{alkyl}$  substituted with cyano;  $\text{C}_{1-6}\text{alkyl}$  substituted with hydroxy and a second substituent selected from cyano or  $\text{R}^7$ ;  $-\text{C}(=\text{N}-\text{O}-\text{R}^8)-\text{C}_{1-4}\text{alkyl}$ ;  $\text{R}^7$  or  $-\text{X}_3-\text{R}^7$ .

-17-

Another interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $R^7$ .

5 Still another interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $C_{1-6}$ alkyl substituted with cyano, in particular  $C_{2-6}$ alkyl substituted with cyano, more in particular ethyl or propyl substituted with cyano; or  $C_{2-6}$ alkenyl substituted with cyano. Preferred is  $C_{2-6}$ alkenyl substituted with cyano.

10 Also an interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $C_{1-6}$ alkyl substituted with cyano and  $R^7$ , or  $C_{2-6}$ alkenyl substituted with cyano and  $R^7$ .

15 A further interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $C_{1-6}$ alkyl substituted with  $R^7$ .

Still a further interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $-C(=N-O-R^8)-C_{1-4}$ alkyl.

20 Also an interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^3$  is  $C_{1-6}$ alkyl substituted with hydroxy and a second substituent selected from cyano or  $R^7$ .

Also an interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $R^2$  or  $R^{2'}$  is cyano or aminocarbonyl and  $R^1$  is hydrogen.

25 Another interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein  $m$  is 2 or 3 and  $X_1$  is  $-NR^5-$ ,  $-O-$ ,  $-C(=O)-$ ,  $-CH_2-$ ,  $-CHOH-$ ,  $-S-$ ,  $-S(=O)_p-$ , in particular wherein  $X_1$  is  $-NR^5-$ , or  $-O-$ .

30 Also an interesting group of compounds are those compounds of formula (I), (I'), (I'') or (I''') wherein one or more, preferably all of the following restrictions apply :

a)  $n$  is at least 1, in particular 1; or  $n'$  is 0;

b)  $R^2$  or  $R^{2'}$  is cyano;

c)  $m$  is 1, 2 or 3;

35 d)  $R^4$  is  $C_{1-6}$ alkyl, especially methyl; nitro; amino; halo;  $C_{1-6}$ alkyloxy or  $R^7$ ;

e)  $R^3$  is  $R^7$ ,  $NR^{13}R^{14}$ ,  $-C(=O)R^{15}$ ,  $-CH=N-NH-C(=O)R^{16}$ ,  $-C(=O)NHR^{13}$ ,

$-C(=O)NR^{13}R^{14}$ ,  $-C(=N-OR^8)-C_{1-4}$ alkyl,  $C_{1-6}$ alkyl substituted with cyano,  $C_{1-6}$ alkyl

substituted twice with cyano, C<sub>1-6</sub>alkyl substituted with NR<sup>9</sup>R<sup>10</sup>, C<sub>1-6</sub>alkyl substituted with hydroxy and cyano, C<sub>1-6</sub>alkyl substituted with hydroxy and R<sup>7</sup>, C<sub>1-6</sub>alkyloxy C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy C<sub>1-6</sub>alkyl substituted with cyano, C<sub>2-6</sub>alkenyl substituted with R<sup>7</sup>, C<sub>2-6</sub>alkenyl substituted with cyano, C<sub>2-6</sub>alkenyl substituted twice with cyano, C<sub>2-6</sub>alkenyl substituted with cyano and R<sup>7</sup>, C<sub>2-6</sub>alkenyl substituted with cyano and -C(=O)-C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl substituted with cyano and halo, C<sub>2-6</sub>alkenyl substituted with -C(=O)-NR<sup>9</sup>R<sup>10</sup>, C<sub>2-6</sub>alkenyl substituted with halo, C<sub>2-6</sub>alkenyl substituted twice with halo or C<sub>2-6</sub>alkenyl substituted with NR<sup>9</sup>R<sup>10</sup>;


f) X<sub>3</sub> is -C(=O)-, -CH<sub>2</sub>-C(=O)-, or -C(=N-OR<sup>8</sup>)-C<sub>1-4</sub>alkanediyl-;

10 g)  $X_1$  is NH or O;

h) R<sup>1</sup> is hydrogen or C<sub>1-4</sub>alkyl.

Preferred compounds of formula (I), (I'), (I'') or (I''') are compounds 1, 25, 84, 133, 152, 179, 233, 239, 247, 248 (see Tables 3, 4 and 5), their *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms thereof.

In general, compounds of formula (I) can be prepared by reacting an intermediate of formula (II) wherein W<sub>1</sub> is a suitable leaving group such as, for example, halo, triflate, tosylate, methylsulfonyl and the like, with an intermediate of formula (III). This reaction can be performed at elevated temperature.




Alternatively, the above reaction can be performed in the presence of a suitable solvent. Suitable solvents are for example acetonitrile, an alcohol, such as for example ethanol, 2-propanol, 2-propanol-HCl; *N,N*-dimethylformamide; *N,N*-dimethylacetamide, 1-methyl-2-pyrrolidinone; 1,4-dioxane, propyleneglycol monomethylether. Preferably the solvent is 2-propanol, 6 N HCl in 2-propanol or acetonitrile, especially acetonitrile. Optionally, sodium hydride may be present.

30 In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies


-19-

generally known in the art such as, for example, extraction, crystallization, distillation, trituration and chromatography.

- Compounds of formula (I) wherein  $R^3$  is  $R^7$  representing a monocyclic, bicyclic or tricyclic aromatic ring system, said  $R^3$  being represented by  $R^{7'}$  and said compounds being represented by formula (I-a), can be prepared by reacting an intermediate of formula (IV) wherein  $W_2$  represents a suitable leaving group such as, for example, halo, hydroxy, triflate, tosylate, thiomethyl, methylsulfonyl, trifluoromethylsulfonyl and the like, with an intermediate of formula (V) wherein  $R^a$  represents a boronate or a tri( $C_{1-4}$ alkyl)stannane, such as tributylstannane, in the presence of a suitable catalyst, such as for example palladium tetrakis(triphenylphosphine), a suitable salt, such as for example disodium carbonate, dipotassium carbonate, and  $Cs_2CO_3$ , and a suitable solvent, such as for example dioxane, dimethyl ether, toluene or an alcohol/water mixture, e.g. MeOH/ $H_2O$ .  $R^a$  may also represent halo, such as for example bromo, in which case the reaction is performed in the presence of 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane.




- Compounds of formula (I) wherein  $R^3$  is  $R^7$  representing a monocyclic, bicyclic or tricyclic saturated ring system, said  $R^3$  being represented by  $R^{7''}$  and said compounds being represented by formula (I-b), can be prepared by reacting an intermediate of formula (IV) with an intermediate of formula (VI).




- Compounds of formula (I) wherein  $R^3$  represents  $C_{1-6}$ alkyl substituted with cyano, said  $R^3$  being represented by  $C_{1-6}$ alkyl-CN and said compounds being represented by formula (I-c), can be prepared by reacting an intermediate of formula (VII) wherein  $W_3$  represents a suitable leaving group, such as for example, halo, e.g. chloro, with a


suitable cyanide salt, such as for example sodium cyanide or potassium cyanide, in the presence of a suitable solvent, such as for example *N,N*-dimethylformamide or dimethylsulfoxide.



- 5 Compounds of formula (I) wherein  $R^3$  represents  $C_{1-6}$ alkyl substituted with  $R^7$ ;  $NR^9R^{10}$  or  $C_{1-6}$ alkyloxy optionally substituted with CN,  $R^7$  or  $NR^9R^{10}$ , said  $R^3$  being represented by  $C_{1-6}$ alkyl-Q wherein Q represents  $R^7$ ;  $NR^9R^{10}$  or  $C_{1-6}$ alkyloxy optionally substituted with CN,  $R^7$  or  $NR^9R^{10}$ , and said compounds being represented by formula (I-d), can be prepared by reacting an intermediate of formula (VII) with an intermediate
- 10 of formula (VIII), optionally in the presence of a suitable salt, such as for example dipotassium carbonate, potassium cyanide, potassium iodide, and a suitable solvent, such as for example acetonitrile.




- 15 Compounds of formula (I) wherein  $R^3$  represents  $-C(=N-O-R^8)-C_{1-4}$ alkyl, said compounds being represented by formula (I-e), can be prepared by reacting an intermediate of formula (IX) with an intermediate of formula (X) in the presence of a suitable solvent, such as an alcohol, e.g. ethanol.




- 20 Compounds of formula (I) wherein  $R^3$  represents  $CR^{c'}=CR^c-CN$  wherein  $R^c$  represents hydrogen or  $C_{1-4}$ alkyl and  $R^{c'}$  represents hydrogen,  $C_{1-4}$ alkyl or  $R^7$ , provided that  $CR^{c'}=CR^c$  is limited to  $C_{2-6}$ alkenyl, said compounds being represented by formula (I-f),

-21-

can be prepared by reacting an intermediate of formula (XI) with a Wittig or Horner-Emmons reagent of formula (XII), wherein  $R^b$ - represents for example  $(\text{Phenyl})_3\text{P}^+-\text{Cl}^-$  or  $(\text{CH}_3\text{CH}_2-\text{O})_2\text{P}(=\text{O})^-$ , which can be considered as a suitable precursor of a phosphorus ylide, in the presence of a suitable salt, such as for example potassium tert.-butoxide, and a suitable solvent, such as for example tetrahydrofuran.



Compounds of formula (I-f-1) and (I-f-2) as depicted below can be prepared by reacting an intermediate of formula (XXXIX) or an appropriate addition salt thereof, wherein  $\text{W}_5$  represents a suitable leaving group, with acrylonitrile or acrylamide in the presence of a suitable palladium catalyst, a suitable base and a suitable solvent.



Suitable leaving groups in the above reaction are for example halo, triflate, tosylate, mesylate and the like. Preferably,  $\text{W}_5$  is halo, more particularly iodo or bromo.

The palladium (Pd) catalyst may be a homogeneous Pd catalyst, such as for example  $\text{Pd}(\text{OAc})_2$ ,  $\text{PdCl}_2$ ,  $\text{Pd}(\text{PPh}_3)_4$ ,  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ , bis(dibenzylidene acetone) palladium, palladium thiomethylphenylglutaramide metallacycle and the like, or a heterogeneous Pd catalyst, such as for example palladium on charcoal, palladium on metal oxides, palladium on zeolites.

-22-

Preferably, the palladium catalyst is a heterogeneous Pd catalyst, more preferably palladium on charcoal (Pd/C). Pd/C is a recoverable catalyst, is stable and relatively inexpensive. It can be easily separated (filtration) from the reaction mixture thereby reducing the risk of Pd traces in the final product. The use of Pd/C also avoids the need for ligands, such as for example phosphine ligands, which are expensive, toxic and contaminants of the synthesized products.


Suitable bases in the above reaction are for example sodium acetate, potassium acetate, *N,N*-diethylethanamine, sodium hydrogencarbonate, sodium hydroxide and the like.

10

Suitable solvents in the above reaction are for example acetonitrile, *N,N*-dimethylacetamide, an ionic liquid e.g. [bmim]PF<sub>6</sub>, *N,N*-dimethylformamide, water, tetrahydrofuran, dimethylsulphoxide, 1-methyl-2-pyrrolidinone and the like.

Compounds of formula (I) wherein R<sup>3</sup> represents CR<sup>c</sup>=CR<sup>c''</sup>-CN with R<sup>c</sup> being as defined hereinabove and R<sup>c''</sup> representing NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>, said compounds being represented by formula (I-g), can be prepared by reacting an intermediate of formula (XI-a) with an intermediate of formula (XIII) in the presence of a suitable solvent, such as for example an alcohol and an alcoholate, e.g. methanol and sodium ethanolate.

20




Compounds of formula (I) wherein R<sup>3</sup> represents CH=C(CN)-CH<sub>2</sub>-CN, said compounds being represented by formula (I-h), can be prepared by reacting an intermediate of formula (XI-b) with 2-butenedinitrile in the presence of tributylphosphine and a suitable solvent, such as for example tetrahydrofuran.


25




-23-



Compounds of formula (I) wherein  $R^3$  represents  $\text{CH}=\text{C}(\text{CN})_2$ , said compounds being represented by formula (I-h'), can be prepared by reacting an intermediate of formula (XI-b) with propanedinitrile in the presence of a suitable base, such as for example piperidine, and a suitable solvent, such as for example an alcohol, e.g. ethanol and the like.




Compounds of formula (I) wherein  $R^3$  represents  $-\text{CHOH}-\text{CH}_2-\text{CN}$ , said compounds being represented by formula (I-i), can be prepared by reacting an intermediate of formula (XI-b) with  $\text{CH}_3-\text{CN}$  in the presence of a suitable proton-abstrating agent, such as for example butyl lithium, in the presence of a suitable substrate for the proton-abstrating agent, for example *N*-(1-methylethyl)-2-propanamine, and in the presence of a suitable solvent, such as for example tetrahydrofuran.




Compounds of formula (I) wherein  $R^3$  represents  $\text{CR}^{\text{c}}=\text{CR}^{\text{c}}-\text{halo}$  wherein  $\text{R}^{\text{c}}$  represents hydrogen or  $\text{C}_{1-4}$ alkyl and  $\text{R}^{\text{c}'}$  represents hydrogen,  $\text{C}_{1-4}$ alkyl or  $\text{R}^7$ , provided that  $\text{CR}^{\text{c}}=\text{CR}^{\text{c}}$  is limited to  $\text{C}_{2-6}$ alkenyl, said compounds being represented by formula (I-j),

-24-

- can be prepared by reacting an intermediate of formula (XI) with a Wittig or Horner-Emmons reagent of formula (XII'), wherein R<sup>b</sup>- represents for example (Phenyl)<sub>3</sub>P<sup>+</sup>- Cl<sup>-</sup> or (CH<sub>3</sub>CH<sub>2</sub>-O)<sub>2</sub>P(=O)-, which can be considered as a suitable precursor of a phosphorus ylide, in the presence of nBuLi, and a suitable solvent, such as for example tetrahydrofuran.




- Compounds of formula (I) wherein R<sup>3</sup> represents CR<sup>c</sup>=CR<sup>c''</sup>-halo with R<sup>c</sup> being as defined hereinabove and R<sup>c''</sup> representing CN, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>, said compounds being represented by formula (I-k), can be prepared by reacting an intermediate of formula (XI-a) with an intermediate of formula (XIII-a) in the presence of a Horner-Emmons reagent such as for example (CH<sub>3</sub>CH<sub>2</sub>-O)<sub>2</sub>P(=O)-Cl, nBuLi, 1,1,1-trimethyl-N-(trimethylsilyl)-silanamine, and a suitable solvent, such as for example tetrahydrofuran.




- Compounds of formula (I) wherein R<sup>3</sup> represents CH=C(Br)<sub>2</sub>, said compounds being represented by formula (I-l), can be prepared by reacting an intermediate of formula (XVIII) with CBr<sub>4</sub>, in the presence of a suitable catalyst salt, such as for example (CuCl)<sub>2</sub>, and in the presence of a suitable base, such as for example NH<sub>3</sub>, and a suitable solvent, such as for example dimethylsulfoxide.


-25-



Compounds of formula (I-m) can be prepared by reacting an intermediate of formula (XIV) with Cl<sub>2</sub>C=S in the presence of a suitable solvent, such as for example dioxane.




- 5 Compounds of formula (I-n) can be prepared by reacting an intermediate of formula (XV) with an intermediate of formula (XVI) in the presence of a suitable solvent, such as for example an alcohol or an alcoholate, e.g. ethanol or sodium methanolate.





- 10 Compounds of formula (I) wherein R<sup>3</sup> represents C<sub>2-6</sub>alkenyl substituted with C(=O)NR<sup>9</sup>R<sup>10</sup> and optionally further substituted with cyano, said compounds being represented by formula (I-o) wherein C<sub>2-6</sub>alkenyl' represents C<sub>2-6</sub>alkenyl optionally substituted with cyano, can be prepared by reacting an intermediate of formula (XXIX) with an intermediate of formula (XXX) in the presence of hydroxybenzotriazole and ethyldimethylaminopropyl carbodiimide and a suitable solvent, such as for example methylene chloride or tetrahydrofuran, and optionally in the presence of a suitable base, such as for example *N,N*-diethylethanamine, NH<sub>4</sub>OH and the like.
- 15

-26-




Compounds of formula (I) wherein R<sup>3</sup> represents  $-\text{C}(=\text{O})\text{NR}^{13}\text{R}^{14}$  or  $-\text{C}(=\text{O})\text{NHR}^{13}$ , said compounds being represented by formula (I-p-1) and (I-p-2) can be prepared by reacting an intermediate of formula (XXXI) with an intermediate of formula (XXXII-1) or (XXXII-2) in the presence of hydroxybenzotriazole and ethyldimethylaminopropyl carbodiimide and a suitable solvent, such as for example methylene chloride or tetrahydrofuran, and optionally in the presence of a suitable base, such as for example *N,N*-diethylethanamine.




- 10 Compounds of formula (I) wherein R<sup>3</sup> represents  $\text{CH}=\text{N}-\text{NH}-\text{C}(=\text{O})-\text{R}^{16}$ , said compounds being represented by formula (I-q), can be prepared by reacting an intermediate of formula (XI-b) with an intermediate of formula (XXXIII) in the presence of a suitable solvent, such as for example methylene chloride and an alcohol, e.g. methanol, ethanol and the like.


-27-



- Compounds of formula (I) wherein  $R^3$  represents  $N(CH_3)_2$ , said compounds being represented by formula (I-r), can be prepared by reductive methylation of an intermediate of formula (XXXIV) with formaldehyde in the presence of a suitable catalyst, such as for example a suitable acid, i.e. acetic acid and the like, palladium on charcoal, Raney Nickel, and in the presence of a suitable reductive agent, such as for example sodium cyanoborohydride or  $H_2$ , and a suitable solvent, such as for example acetonitrile.




- Compounds of formula (I) wherein  $R^3$  represents pyrrolyl, said compounds being represented by formula (I-s), can be prepared by reacting an intermediate of formula (XXXIV) with 2,5-dimethoxytetrahydrofuran in the presence of a suitable acid, such as for example acetic acid.



- Compounds of formula (I) wherein  $R^3$  represents  $CH=CH-R^7$ , said compounds being represented by formula (I-t), can be prepared by reacting an intermediate of formula (XXXV) (Ph indicates phenyl) with an intermediate of formula (XXXVI) in the presence of  $nBuLi$  and a suitable solvent, such as for example tetrahydrofuran.

-28-



The compounds of formula (I) may further be prepared by converting compounds of formula (I) into each other according to art-known group transformation reactions.

- 5 The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth
- 10 alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxy-peroxy acid or halo substituted benzenecarboxy-peroxy acid, e.g. 3-chlorobenzenecarboxy-peroxy acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. tert.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols,
- 15 e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

For instance, a compound of formula (I) wherein  $R^3$  comprises cyano, can be converted into a compound of formula (I) wherein  $R^3$  comprises aminocarbonyl, by reaction with

20  $HCOOH$ , in the presence of a suitable acid, such as hydrochloric acid. A compound of formula (I) wherein  $R^3$  comprises cyano, can also further be converted into a compound of formula (I) wherein  $R^3$  comprises tetrazolyl, by reaction with sodium azide in the presence of ammonium chloride and *N,N*-dimethylacetamide.

- 25 Compounds of formula (I) wherein  $R^3$  comprises aminocarbonyl, can be converted into a compound of formula (I) wherein  $R^3$  comprises cyano, in the presence of a suitable dehydrating agent. The dehydration can be performed according to methodologies well-known to the person skilled in the art, such as the ones disclosed in "Comprehensive Organic Transformations. A guide to functional group preparations"
- 30 by Richard C. Larock, John Wiley & Sons, Inc, 1999, p 1983-1985, which is

-29-

incorporated herein as reference. Different suitable reagents are enumerated in said reference, such as for example  $\text{SOCl}_2$ ,  $\text{HOSO}_2\text{NH}_2$ ,  $\text{ClSO}_2\text{NCO}$ ,  $\text{MeO}_2\text{CNSO}_2\text{NEt}_3$ ,  $\text{PhSO}_2\text{Cl}$ ,  $\text{TsCl}$ ,  $\text{P}_2\text{O}_5$ ,  $(\text{Ph}_3\text{PO}_3\text{SCF}_3)_3\text{O}_3\text{SCF}_3$ , polyphosphate ester,  $(\text{EtO})_2\text{POP}(\text{OEt})_2$ ,  $(\text{EtO})_3\text{PI}_2$ , 2-chloro-1,3,2-dioxaphospholane, 2,2,2-trichloro-2,2-dihydro-1,3,2-dioxaphospholane,  $\text{POCl}_3$ ,  $\text{PPh}_3$ ,  $\text{P}(\text{NCl}_2)_3$ ,  $\text{P}(\text{NEt}_2)_3$ ,  $\text{COCl}_2$ ,  $\text{NaCl}.\text{AlCl}_3$ ,  $\text{ClCOCOCl}$ ,  $\text{ClCO}_2\text{Me}$ ,  $\text{Cl}_3\text{CCOCl}$ ,  $(\text{CF}_3\text{CO})_2\text{O}$ ,  $\text{Cl}_3\text{CN}=\text{CCl}_2$ , 2,4,6-trichloro-1,3,5-triazine,  $\text{NaCl}.\text{AlCl}_3$ ,  $\text{HN}(\text{SiMe}_2)_3$ ,  $\text{N}(\text{SiMe}_2)_4$ ,  $\text{LiAlH}_4$  and the like. All the reagents listed in said publication are incorporated herein as reference.

10 Compounds of formula (I) wherein  $\text{R}^3$  comprises  $\text{C}_{2-6}$ alkenyl can be converted into a compound of formula (I) wherein  $\text{R}^3$  comprises  $\text{C}_{1-6}$ alkyl by reduction in the presence of a suitable reducing agent, such as for example  $\text{H}_2$ , in the presence of a suitable catalyst, such as for example palladium on charcoal, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.

15 Compounds of formula (I) wherein  $\text{R}^3$  represents  $\text{CH}(\text{OH})-\text{R}^{16}$ , can be converted into a compound of formula (I) wherein  $\text{R}^3$  represents  $\text{C}(=\text{O})-\text{R}^{16}$  by reaction with Jones's reagent in the presence of a suitable solvent, such as for example 2-propanone.

20 Compound of formula (I) wherein  $\text{R}^3$  represents  $\text{C}(=\text{O})-\text{CH}_2-\text{R}^{16a}$ , wherein  $\text{R}^{16a}$  represents cyano or aminocarbonyl, can be converted into a compound of formula (I) wherein  $\text{R}^3$  represents  $\text{C}(\text{Cl})=\text{CH}-\text{R}^{16a}$  by reaction with  $\text{POCl}_3$ .

25 Compounds of formula (I) wherein  $\text{R}^3$  represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with formyl can be converted into compounds of formula (I) wherein  $\text{R}^3$  represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with  $\text{CH}(=\text{N}-\text{O}-\text{R}^8)$  by reaction with  $\text{NH}_2\text{OR}^8$  in the presence of a suitable base, such as for example sodium hydroxide and a suitable solvent, such as for example an alcohol, e.g. ethanol and the like. Compounds of formula (I) wherein  $\text{R}^3$  represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with  $\text{CH}(=\text{N}-\text{O}-\text{R}^8)$  can be converted into a compound of formula (I) wherein  $\text{R}^3$  represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated

-30-

or aromatic heterocycle substituted with CN by reaction with a carbodiimide in the presence of a suitable solvent, such as for example tetrahydrofuran.


- 5 Compounds of formula (I) wherein  $R^4$  represents nitro, can be converted into a compound of formula (I) wherein  $R^4$  is amino, in the presence of a suitable reducing agent, such as for example  $H_2$ , in the presence of a suitable catalyst, such as for example Raney Nickel, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.
- 10 Compounds of formula (I) wherein  $R^1$  is hydrogen, can be converted into a compound of formula (I) wherein  $R^1$  is  $C_{1-6}$ alkyl, by reaction with a suitable alkylating agent, such as for example iodo- $C_{1-6}$ alkyl, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example tetrahydrofuran.
- 15 Some of the compounds of formula (I) and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter
- 20 current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques,
- 25 e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereochemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.
- 30 An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.
- 35 Some of the intermediates and starting materials are known compounds and may be




-31-

commercially available or may be prepared according to art-known procedures or some of the compounds of formula (I) or the described intermediates may be prepared according to the procedures described in WO 99/50250 and WO 00/27825.


- 5 Intermediates of formula (II) can be prepared by reacting an intermediate of formula (XVII) with a leaving group introducing agent of formula (XIX) wherein  $W_1$  represents the leaving group and R represents the remaining of the leaving group introducing agent, such as for example  $POCl_3$ .



- 10 Intermediates of formula (III) wherein  $X_1$  represents NH, said intermediates being represented by formula (III-a), can be prepared from an intermediate of formula (XX) in the presence of  $ZnCl_2$  and in the presence of a suitable solvent, such as for example an alcohol, for example ethanol.




- 15 Intermediates of formula (III'-a) as depicted below can be prepared from an intermediate of formula (XX) wherein  $R^3$  represents  $C_{2-6}$ alkenyl substituted with CN, said intermediate being represented by formula (XX-a), in the presence of  $ZnCl_2$  and in the presence of a suitable  $C_{1-4}$ alkyl-OH, such as for example ethanol.



- 20 Intermediates of formula (III-b-1) and (III-b-2) as depicted below can be prepared by reacting an intermediate of formula (XLI) or an appropriate acid addition salt thereof,

-32-

wherein  $W_6$  represents a suitable leaving group, with acrylonitrile or acrylamide in the presence of a suitable palladium catalyst, a suitable base and a suitable solvent.



- 5 Suitable leaving groups in the above reaction are for example halo, triflate, tosylate, mesylate and the like. Preferably,  $W_6$  is halo, more preferably iodo or bromo.

- The palladium (Pd) catalyst may be a homogeneous Pd catalyst, such as for example  $Pd(OAc)_2$ ,  $PdCl_2$ ,  $Pd(PPh_3)_4$ ,  $Pd(PPh_3)_2Cl_2$ , bis(dibenzylidene acetone) palladium, palladium thiomethylphenylglutaramide metallacycle and the like, or a heterogeneous Pd catalyst, such as for example palladium on charcoal, palladium on metal oxides, palladium on zeolites.

- 10 Preferably, the palladium catalyst is a heterogeneous Pd catalyst, more preferably palladium on charcoal (Pd/C). Pd/C is a recoverable catalyst, is stable and relatively inexpensive. It can be easily separated (filtration) from the reaction mixture thereby reducing the risk of Pd traces in the final product. The use of Pd/C also avoids the need for ligands, such as for example phosphine ligands, which are expensive, toxic and contaminants of the synthesized products.


- 20 Suitable bases in the above reaction are for example sodium acetate, potassium acetate, *N,N*-diethylethanamine, sodium hydrogencarbonate, sodium hydroxide and the like.

- Suitable solvents in the above reaction are for example acetonitrile, *N,N*-dimethylacetamide, an ionic liquid e.g. [bmim]PF<sub>6</sub>, *N,N*-dimethylformamide, water, tetrahydrofuran, dimethylsulphoxide, 1-methyl-2-pyrrolidinone and the like.


- 25

-33-

- Intermediates of formula (III-b-2) can be converted into an intermediate of formula (III-b-1) in the presence of a suitable dehydrating agent. The dehydration can be performed according to methodologies well-known to the person skilled in the art, such as the ones disclosed in "Comprehensive Organic Transformations. A guide to functional group preparations" by Richard C. Larock, John Wiley & Sons, Inc, 1999, p 1983-1985, which is incorporated herein as reference. Different suitable reagents are enumerated in said reference, such as for example  $\text{SOCl}_2$ ,  $\text{HOSO}_2\text{NH}_2$ ,  $\text{ClSO}_2\text{NCO}$ ,  $\text{MeO}_2\text{CNSO}_2\text{NEt}_3$ ,  $\text{PhSO}_2\text{Cl}$ ,  $\text{TsCl}$ ,  $\text{P}_2\text{O}_5$ ,  $(\text{Ph}_3\text{PO}_3\text{SCF}_3)_3\text{O}_3\text{SCF}_3$ , polyphosphate ester,  $(\text{EtO})_2\text{POP}(\text{OEt})_2$ ,  $(\text{EtO})_3\text{PI}_2$ , 2-chloro-1,3,2-dioxaphospholane, 2,2,2-trichloro-2,2-dihydro-1,3,2-dioxaphospholane,  $\text{POCl}_3$ ,  $\text{PPh}_3$ ,  $\text{P}(\text{NCl}_2)_3$ ,  $\text{P}(\text{NEt}_2)_3$ ,  $\text{COCl}_2$ ,  $\text{NaCl}.\text{AlCl}_3$ ,  $\text{ClCOCOCOC}$ ,  $\text{ClCO}_2\text{Me}$ ,  $\text{Cl}_3\text{CCOCl}$ ,  $(\text{CF}_3\text{CO})_2\text{O}$ ,  $\text{Cl}_3\text{CN}=\text{CCl}_2$ , 2,4,6-trichloro-1,3,5-triazine,  $\text{NaCl}.\text{AlCl}_3$ ,  $\text{HN}(\text{SiMe}_2)_3$ ,  $\text{N}(\text{SiMe}_2)_4$ ,  $\text{LiAlH}_4$  and the like. All the reagents listed in said publication are incorporated herein as reference.
- Intermediates of formula (XX) wherein  $\text{R}^3$  represents  $\text{CR}^c=\text{CR}^c-\text{CN}$  with  $\text{R}^c$  and  $\text{R}^{c'}$  as described hereinabove, said intermediates being represented by formula (XX-b), can be prepared from an intermediate of formula (XXI) by the reaction described above for the preparation of a compound of formula (I-f).




- Intermediates of formula (XXI) can be prepared by oxidation of an intermediate of formula (XXII) in the presence of a suitable oxidizing agent, such as for example  $\text{KMnO}_4$ , in the presence of a suitable solvent, such as for example methylene chloride and tris[2-(2-methoxyethoxy)ethyl]amine.




- Intermediates of formula (XXI) wherein  $\text{R}^{c'}$  is H, said intermediates being represented by formula (XXI-a), can also be prepared by reacting an intermediate of formula

-34-


(XXIII) wherein  $W_4$  represents a suitable leaving group, such as halo, e.g. bromo, with *N,N*-dimethylformamide in the presence of *n*BuLi and in the presence of a suitable solvent, such as for example tetrahydrofuran.



- 5 Intermediates of formula (XXII) wherein  $R^{c'}$  represents  $C_{1-4}$ alkyl, said intermediates being represented by formula (XXII-a), can be prepared by reacting an intermediate of formula (XXIII) with an intermediate of formula (XXIV) in the presence of *n*BuLi and in the presence of a suitable solvent, such as for example tetrahydrofuran.




- 10 Intermediates of formula (XI) can be prepared by reacting an intermediate of formula (XXV) with an intermediate of formula (II), optionally in the presence of a suitable base, such as for example 1-methyl-pyrrolidin-2-one, or a suitable acid, such as for example hydrochloric acid.




- 15 Intermediates of formula (XV) can be prepared by reacting an intermediate of formula (XXVI) with an intermediate of formula (II) in the presence of a suitable base, such as for example 1-methyl-pyrrolidin-2-one and sodium hydride and a suitable solvent, such as for example dioxane.

-35-




Intermediates of formula (VII) can be prepared by reacting an intermediate of formula (XXVII) with a leaving group introducing agent of formula (XIX'), such as for example  $\text{SOCl}_2$ , in the presence of a suitable solvent, such as for example methylene chloride.



5


Intermediates of formula (XXVII) wherein  $\text{C}_{1-6}\text{alkyl}$  represents  $\text{CH}_2$ , said intermediates being represented by formula (XXVII-a), can be prepared by reducing an intermediate of formula (XV) or formula (XXXI) with a suitable reducing agent, such as for example  $\text{LiAlH}_4$ , in the presence of a suitable solvent, such as for example tetrahydrofuran.



10


Intermediates of formula (XXVII-a) can be converted to an intermediate of formula (XXXI) by reaction with Jones reagent in the presence of a suitable solvent, such as for example acetone.

-36-




Intermediates of formula (XI-b) can be prepared by oxidizing an intermediate of formula (XXVII-a) in the presence of a suitable oxidizing agent, such as for example  $MnO_2$ , and a suitable solvent, such as for example methylene chloride, *N,N*-dimethylformamide.

5





Intermediates of formula (XIV) can be prepared by reacting an intermediate of formula (XV) with  $H_2N-NH_2$  in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol and the like.




10

Intermediates of formula (IX) and (XI-a) can be reduced to an intermediate of formula (XXVII'-a) and (XXVII'-b) in the presence of a suitable reducing agent, such as for example  $NaBH_4$ ,  $LiAlH_4$  or  $BuLi$  and a suitable solvent, such as for example tetrahydrofuran or an alcohol, e.g. methanol, ethanol and the like.

-37-




An intermediate of formula (XI-b) can be converted into an intermediate of formula (XXVII'-a) by reaction with C<sub>1-4</sub>alkyl-Iodide in the presence of Mg and a suitable solvent, such as for example diethylether and tetrahydrofuran.




5

Intermediates of formula (XVIII) can be prepared by reacting an intermediate of formula (XI-b) with H<sub>2</sub>N-NH<sub>2</sub> in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol and the like.




- 10 Intermediates of formula (XXIX) or (XXXI) can be prepared by hydrolizing an intermediate of formula (XXXVII) wherein C<sub>2-6</sub>alkenyl' represents C<sub>2-6</sub>alkenyl optionally substituted cyano, or an intermediate of formula (XV) in the presence of a suitable aqueous acid solution, such as for example hydrochloric acid 2N and the like, and in the presence of a suitable solvent, such as for example an alcohol, e.g.
- 15 isopropanol and the like.

-38-




- Intermediates of formula (XXXVII) wherein  $\text{C}_{2-6}\text{alkenyl}'$  is  $\text{CH}=\text{CH}$ , said intermediates being represented by formula (XXXVII-a), can be prepared by reacting an intermediate of formula (XI-b) with a Wittig or Horner-Emmons reagent of formula (XII'), wherein  $\text{R}^b$  represents for example  $(\text{Phenyl})_3\text{P}^+-\text{Cl}^-$  or  $(\text{CH}_3\text{CH}_2-\text{O})_2\text{P}(=\text{O})^-$ , which can be considered as a suitable precursor of a phosphorus ylide, in the presence of a suitable solvent, such as for example tetrahydrofuran.




- Intermediates of formula (XXXVII) wherein  $\text{C}_{2-6}\text{alkenyl}'$  is  $-\text{CH}=\text{C}(\text{CN})-$ , said intermediates being represented by formula (XXXVII-b), can be prepared by reacting an intermediate of formula (XI-b) with  $\text{NC}-\text{CH}_2-\text{C}(=\text{O})\text{O}-\text{C}_{1-6}\text{alkyl}$ , in the presence of a suitable base, such as for example piperidine, and a suitable solvent, such as for example an alcohol, e.g. ethanol.




-39-



Intermediates of formula (XXXIV) can be prepared by reducing an intermediate of formula (XXXVIII) in the presence H<sub>2</sub> and a suitable catalyst, such as for example palladium on charcoal or Raney Nickel, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol and the like.




Intermediates of formula (XXXV) can be prepared by reacting an intermediate of formula (VII-a) in the presence of triphenylphosphine and a suitable solvent, such as for example acetonitrile.



Intermediates of formula (XXXIX) can be prepared by reacting an intermediate of formula (XL) with an intermediate of formula (II-a) wherein W<sub>5</sub> and W<sub>1</sub> are as defined hereinbefore.

-40-



The compounds of formula (I) as prepared in the hereinabove described processes may be synthesized as a mixture of stereoisomeric forms, in particular in the form of racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

It will be appreciated by those skilled in the art that in the processes described above the functional groups of intermediate compounds may need to be blocked by protecting groups.

Functional groups which it is desirable to protect include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), benzyl and tetrahydropyranyl. Suitable protecting groups for amino include *tert*-butoxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C<sub>1-6</sub>alkyl or benzyl esters.


The protection and deprotection of functional groups may take place before or after a reaction step.

-41-

The use of protecting groups is fully described in 'Protective Groups in Organic Chemistry', edited by J W F McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis' 2<sup>nd</sup> edition, T W Greene & P G M Wutz, Wiley Interscience (1991).


- 5 The present invention also concerns new compounds of formula (VII), (XXVII), (XXIX) and (XXXVII) which can be used as intermediates in the synthesis of the compounds of formula (I) and which also exhibit HIV replication inhibiting activity.

In particular, the present invention also relates to a compound of formula




- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined hereinabove for the compounds of formula (I) and  $W_3$  represents a suitable leaving group such as for
- 15 example halo, e.g. chloro and the like.

The present invention also relates to a compound of formula



- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined hereinabove for the compounds of formula (I).

The present invention also relates to a compound of formula




-42-

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined hereinabove for the compounds of formula (I) and  $C_{2-6}alkenyl'$  represents  $C_{2-6}alkenyl$  optionally substituted with cyano.

The present invention also relates to a compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined hereinabove for the compounds of formula (I) and  $C_{2-6}alkenyl'$  represents  $C_{2-6}alkenyl$  optionally substituted with cyano.

Compounds of formula (III-b) as depicted below intervene in the synthesis of compounds of formula (I).

Therefore, the present invention also relates to a compound of formula (III-b)



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^4$  and  $X_1$  are as defined hereinabove for the compounds of formula (I).

Preferred compounds of formula (III-b) are those compounds wherein  $X_1$  represents

NH. More preferred compounds of formula (III-b) are those compounds wherein  $X_1$

-43-

represents NH and C<sub>2-6</sub>alkenyl represents CH=CH. Most preferred compounds of formula (III-b) are the compounds of formula (III-b-1) as described hereinabove.

5 The compounds of formula (I), (I'), (I''), (I'''), (VII), (XXVII), (XXIX) and (XXXVII) show antiretroviral properties (reverse transcriptase inhibiting properties), in particular against Human Immunodeficiency Virus (HIV), which is the aetiological agent of Acquired Immune Deficiency Syndrome (AIDS) in humans. The HIV virus preferentially infects human T-4 cells and destroys them or changes their normal function, particularly the coordination of the immune system. As a result, an infected patient has an ever decreasing number of T-4 cells, which moreover behave abnormally. Hence, the immunological defense system is unable to combat infections and neoplasms and the HIV infected subject usually dies by opportunistic infections such as pneumonia, or by cancers. Other conditions associated with HIV infection include thrombocytopaenia, Kaposi's sarcoma and infection of the central nervous system characterized by progressive demyelination, resulting in dementia and symptoms such as, progressive dysarthria, ataxia and disorientation. HIV infection further has also been associated with peripheral neuropathy, progressive generalized lymphadenopathy (PGL) and AIDS-related complex (ARC).

20 The present compounds also show activity against (multi) drug resistant HIV strains, in particular (multi) drug resistant HIV-1 strains, more in particular the present compounds show activity against HIV strains, especially HIV-1 strains, that have acquired resistance to one or more art-known non-nucleoside reverse transcriptase inhibitors. Art-known non-nucleoside reverse transcriptase inhibitors are those non-nucleoside reverse transcriptase inhibitors other than the present compounds and in particular commercial non-nucleoside reverse transcriptase inhibitors. The present compounds also have little or no binding affinity to human  $\alpha$ -1 acid glycoprotein; human  $\alpha$ -1 acid glycoprotein does not or only weakly affect the anti HIV activity of the present compounds.

30 Due to their antiretroviral properties, particularly their anti-HIV properties, especially their anti-HIV-1-activity, the compounds of formula (I), their *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms thereof, are useful in the treatment of individuals infected by HIV and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the enzyme reverse

-44-

transcriptase. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic Central Nervous System diseases caused  
5 by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

The compounds of the present invention or any subgroup thereof may therefore be used as medicines against above-mentioned conditions. Said use as a medicine or method of treatment comprises the administration to HIV-infected subjects of an amount effective  
10 to combat the conditions associated with HIV and other pathogenic retroviruses, especially HIV-1. In particular, the compounds of formula (I) may be used in the manufacture of a medicament for the treatment or the prevention of HIV infections.

In view of the utility of the compounds of formula (I), there is provided a method of  
15 treating warm-blooded animals, including humans, suffering from or a method of preventing warm-blooded animals, including humans, to suffer from viral infections, especially HIV infections. Said method comprises the administration, preferably oral administration, of an effective amount of a compound of formula (I), a *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine or a possible  
20 stereoisomeric form thereof, to warm-blooded animals, including humans.

The present invention also provides compositions for treating viral infections comprising a therapeutically effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier or diluent.

25 The compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an  
30 effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally,  
35 rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of

-45-

oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most

5 advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and

10 glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a

15 suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

20 The compounds of the present invention may also be administered via inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder. Any system developed for the delivery of solutions, suspensions or dry

25 powders via oral or nasal inhalation or insufflation are suitable for the administration of the present compounds.

To aid solubility of the compounds of formula (I), suitable ingredients, e.g. cyclodextrins, may be included in the compositions. Appropriate cyclodextrins are  $\alpha$ -,

30  $\beta$ -,  $\gamma$ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with  $C_{1-6}$ alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated  $\beta$ -CD; hydroxy $C_{1-6}$ alkyl, particularly hydroxyethyl, hydroxy-propyl or hydroxybutyl; carboxy $C_{1-6}$ alkyl, particularly carboxymethyl or carboxy-ethyl;  $C_{1-6}$ alkylcarbonyl,

35 particularly acetyl. Especially noteworthy as complexants and/or solubilizers are  $\beta$ -CD, randomly methylated  $\beta$ -CD, 2,6-dimethyl- $\beta$ -CD, 2-hydroxyethyl- $\beta$ -CD,

-46-

2-hydroxyethyl- $\beta$ -CD, 2-hydroxypropyl- $\beta$ -CD and (2-carboxymethoxy)propyl- $\beta$ -CD, and in particular 2-hydroxypropyl- $\beta$ -CD (2-HP- $\beta$ -CD).

5 The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The M.S. and D.S. value can be determined by various analytical techniques such as nuclear magnetic resonance (NMR), mass spectrometry (MS) and infrared spectroscopy (IR). Depending on the technique used, slightly different values may be obtained for one given cyclodextrin derivative. Preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10 and the D.S. ranges from 0.125 to 3.

Other suitable compositions for oral or rectal administration comprise particles consisting of a solid dispersion comprising a compound of formula (I) and one or more appropriate pharmaceutically acceptable water-soluble polymers.

20 The term "a solid dispersion" used hereinafter defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, in casu the compound of formula (I) and the water-soluble polymer, wherein one component is dispersed more or less evenly throughout the other component or components ( in case additional pharmaceutically acceptable formulating agents, generally known in the art, are included, such as plasticizers, preservatives and the like). When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermo-dynamics, such a solid dispersion will be called "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered. This advantage can probably be explained by the ease with which said solid solutions can form liquid solutions when contacted with a liquid medium such as the gastro-intestinal juices. The ease of dissolution may be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of components from a crystalline or microcrystalline solid phase.



The term "a solid dispersion" also comprises dispersions which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase. For example, the term "a solid dispersion" also relates to a system having domains or small regions wherein  
5 amorphous, microcrystalline or crystalline compound of formula (I), or amorphous, microcrystalline or crystalline water-soluble polymer, or both, are dispersed more or less evenly in another phase comprising water-soluble polymer, or compound of formula (I), or a solid solution comprising compound of formula (I) and water-soluble polymer. Said domains are regions within the solid dispersion distinctively marked by  
10 some physical feature, small in size, and evenly and randomly distributed throughout the solid dispersion.

Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation.

15

The solution-evaporation process comprises the following steps :

- a) dissolving the compound of formula (I) and the water-soluble polymer in an appropriate solvent, optionally at elevated temperatures;
- b) heating the solution resulting under point a), optionally under vacuum, until the  
20 solvent is evaporated. The solution may also be poured onto a large surface so as to form a thin film, and evaporating the solvent therefrom.

In the spray-drying technique, the two components are also dissolved in an appropriate solvent and the resulting solution is then sprayed through the nozzle of a spray dryer  
25 followed by evaporating the solvent from the resulting droplets at elevated temperatures.

The preferred technique for preparing solid dispersions is the melt-extrusion process comprising the following steps :

- 30 a) mixing a compound of formula (I) and an appropriate water-soluble polymer,
- b) optionally blending additives with the thus obtained mixture,
- c) heating and compounding the thus obtained blend until one obtains a homogenous melt,
- d) forcing the thus obtained melt through one or more nozzles; and
- 35 e) cooling the melt till it solidifies.

-48-

The terms "melt" and "melting" should be interpreted broadly. These terms not only mean the alteration from a solid state to a liquid state, but can also refer to a transition to a glassy state or a rubbery state, and in which it is possible for one component of the mixture to get embedded more or less homogeneously into the other. In particular cases, one component will melt and the other component(s) will dissolve in the melt thus forming a solution, which upon cooling may form a solid solution having advantageous dissolution properties.

After preparing the solid dispersions as described hereinabove, the obtained products can be optionally milled and sieved.

The solid dispersion product may be milled or ground to particles having a particle size of less than 600  $\mu\text{m}$ , preferably less than 400  $\mu\text{m}$  and most preferably less than 125  $\mu\text{m}$ .

The particles prepared as described hereinabove can then be formulated by conventional techniques into pharmaceutical dosage forms such as tablets and capsules.

It will be appreciated that a person of skill in the art will be able to optimize the parameters of the solid dispersion preparation techniques described above, such as the most appropriate solvent, the working temperature, the kind of apparatus being used, the rate of spray-drying, the throughput rate in the melt-extruder

The water-soluble polymers in the particles are polymers that have an apparent viscosity, when dissolved at 20°C in an aqueous solution at 2 % (w/v), of 1 to 5000 mPa.s more preferably of 1 to 700 mPa.s, and most preferred of 1 to 100 mPa.s. For example, suitable water-soluble polymers include alkylcelluloses, hydroxyalkylcelluloses, hydroxyalkyl alkylcelluloses, carboxyalkylcelluloses, alkali metal salts of carboxyalkylcelluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters, starches, pectines, chitin derivatives, di-, oligo- and polysaccharides such as trehalose, alginic acid or alkali metal and ammonium salts thereof, carrageenans, galactomannans, tragacanth, agar-agar, gummi arabicum, guar gummi and xanthan gummi, polyacrylic acids and the salts thereof, polymethacrylic acids and the salts thereof, methacrylate copolymers, polyvinylalcohol, polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate, combinations of polyvinylalcohol and polyvinylpyrrolidone, polyalkylene oxides and copolymers of ethylene oxide and propylene oxide. Preferred water-soluble polymers are hydroxypropyl methylcelluloses.

Also one or more cyclodextrins can be used as water soluble polymer in the preparation of the above-mentioned particles as is disclosed in WO 97/18839. Said cyclodextrins include the pharmaceutically acceptable unsubstituted and substituted cyclodextrins known in the art, more particularly  $\alpha$ ,  $\beta$  or  $\gamma$  cyclodextrins or the pharmaceutically acceptable derivatives thereof.

Substituted cyclodextrins which can be used to prepare the above described particles include polyethers described in U.S. Patent 3,459,731. Further substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C<sub>1</sub>-6alkyl, hydroxyC<sub>1</sub>-6alkyl, carboxy-C<sub>1</sub>-6alkyl or C<sub>1</sub>-6alkyloxycarbonylC<sub>1</sub>-6alkyl or mixed ethers thereof. In particular such substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C<sub>1</sub>-3alkyl, hydroxyC<sub>2</sub>-4alkyl or carboxyC<sub>1</sub>-2alkyl or more in particular by methyl, ethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, carboxymethyl or carboxyethyl.

Of particular utility are the  $\beta$ -cyclodextrin ethers, e.g. dimethyl- $\beta$ -cyclodextrin as described in Drugs of the Future, Vol. 9, No. 8, p. 577-578 by M. Nogradi (1984) and polyethers, e.g. hydroxypropyl  $\beta$ -cyclodextrin and hydroxyethyl  $\beta$ -cyclodextrin, being examples. Such an alkyl ether may be a methyl ether with a degree of substitution of about 0.125 to 3, e.g. about 0.3 to 2. Such a hydroxypropyl cyclodextrin may for example be formed from the reaction between  $\beta$ -cyclodextrin and propylene oxide and may have a MS value of about 0.125 to 10, e.g. about 0.3 to 3.

Another type of substituted cyclodextrins is sulfobutylcyclodextrines.

The ratio of the compound of formula (I) over the water soluble polymer may vary widely. For example ratios of 1/100 to 100/1 may be applied. Interesting ratios of the compound of formula (I) over cyclodextrin range from about 1/10 to 10/1. More interesting ratios range from about 1/5 to 5/1.

It may further be convenient to formulate the compounds of formula (I) in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those which physically adhere to the surface of the compound of formula (I) but do not chemically bond to said compound.

-50-

Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

Yet another interesting way of formulating the compounds of formula (I) involves a pharmaceutical composition whereby the compounds of formula (I) are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.

Said beads comprise a central, rounded or spherical core, a coating film of a hydrophilic polymer and a compound of formula (I) and optionally a seal-coating layer.

Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof.

Those of skill in the treatment of HIV-infection could determine the effective daily amount from the test results presented here. In general it is contemplated that an effective daily amount would be from 0.01 mg/kg to 50 mg/kg body weight, more preferably from 0.1 mg/kg to 10 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

-51-

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines and are not intended to limit the scope or use of the invention to any extent.

The present compounds of formula (I) can be used alone or in combination with other therapeutic agents, such as anti-virals, antibiotics, immunomodulators or vaccines for the treatment of viral infections. They may also be used alone or in combination with other prophylactic agents for the prevention of viral infections. The present compounds may be used in vaccines and methods for protecting individuals against viral infections over an extended period of time. The compounds may be employed in such vaccines either alone or together with other compounds of this invention or together with other anti-viral agents in a manner consistent with the conventional utilization of reverse transcriptase inhibitors in vaccines. Thus, the present compounds may be combined with pharmaceutically acceptable adjuvants conventionally employed in vaccines and administered in prophylactically effective amounts to protect individuals over an extended period of time against HIV infection.

Also, the combination of an antiretroviral compound and a compound of formula (I) can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of formula (I), and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in anti-HIV treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. Said other antiretroviral compounds may be known antiretroviral compounds such as suramine, pentamidine, thymopentin, castanospermine, dextran (dextran sulfate), foscarnet-sodium (trisodium phosphono formate); nucleoside reverse transcriptase inhibitors, e.g. zidovudine (3'-azido-3'-deoxythymidine, AZT), didanosine (2',3'-dideoxyinosine; ddI), zalcitabine (dideoxycytidine, ddC) or lamivudine (2'-3'-dideoxy-3'-thiacytidine, 3TC), stavudine (2',3'-didehydro-3'-deoxythymidine, d4T), abacavir and the like; non-nucleoside reverse transcriptase inhibitors such as nevirapine (11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido-

-52-

- [3,2-b : 2',3'-e][1,4]diazepin-6-one), efavirenz, delavirdine, TMC-120, TMC-125 and the like; phosphonate reverse transcriptase inhibitors, e.g. tenofovir and the like; compounds of the TIBO (tetrahydro-imidazo[4,5,1-jk][1,4]-benzodiazepine-2(1*H*)-one and thione)-type e.g. (S)-8-chloro-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-2-butenyl)-imidazo-[4,5,1-jk][1,4]benzodiazepine-2(1*H*)-thione; compounds of the  $\alpha$ -APA ( $\alpha$ -anilino phenyl acetamide) type e.g.  $\alpha$ -[(2-nitrophenyl)amino]-2,6-dichlorobenzene-acetamide and the like; inhibitors of trans-activating proteins, such as TAT-inhibitors, e.g. RO-5-3335, or REV inhibitors, and the like; protease inhibitors e.g. indinavir, ritonavir, saquinavir, lopinavir (ABT-378), nelfinavir, amprenavir, TMC-126, BMS-232632, VX-175 and the like; fusion inhibitors, e.g. T-20, T-1249 and the like; CXCR4 receptor antagonists, e.g. AMD-3100 and the like; inhibitors of the viral integrase; nucleotide-like reverse transcriptase inhibitors, e.g. tenofovir and the like; ribonucleotide reductase inhibitors, e.g. hydroxyurea and the like.
- By administering the compounds of the present invention with other anti-viral agents which target different events in the viral life cycle, the therapeutic effect of these compounds can be potentiated. Combination therapies as described above exert a synergistic effect in inhibiting HIV replication because each component of the combination acts on a different site of HIV replication. The use of such combinations may reduce the dosage of a given conventional anti-retroviral agent which would be required for a desired therapeutic or prophylactic effect as compared to when that agent is administered as a monotherapy. These combinations may reduce or eliminate the side effects of conventional single anti-retroviral therapy while not interfering with the anti-viral activity of the agents. These combinations reduce potential of resistance to single agent therapies, while minimizing any associated toxicity. These combinations may also increase the efficacy of the conventional agent without increasing the associated toxicity.
- The compounds of the present invention may also be administered in combination with immunomodulating agents, e.g. levamisole, bropirimine, anti-human alpha interferon antibody, interferon alpha, interleukin 2, methionine enkephalin, diethyldithiocarbamate, tumor necrosis factor, naltrexone and the like; antibiotics, e.g. pentamidine isethionate and the like; cholinergic agents, e.g. tacrine, rivastigmine, donepezil, galantamine and the like; NMDA channel blockers, e.g. memantine to prevent or combat infection and diseases or symptoms of diseases associated with HIV infections, such as AIDS and ARC, e.g. dementia. A compound of formula (I) can also be combined with another compound of formula (I).

Although the present invention focuses on the use of the present compounds for preventing or treating HIV infections, the present compounds may also be used as inhibitory agents for other viruses which depend on similar reverse transcriptases for obligatory events in their life cycle.

The following examples are intended to illustrate the present invention.


#### Experimental part

Hereinafter, "DMF" is defined as *N,N*-dimethylformamide, "DIPE" is defined as diisopropyl ether, "THF" is defined as tetrahydrofuran, "DMA" is defined as *N,N*-dimethylacetamide, "DMSO" is defined as dimethylsulfoxide, "DME" is defined as dimethyl ether, "EtOAc" is defined as ethylacetate, "EDCI" is defined as *N*'-(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine.

#### A. Preparation of the intermediate compounds

##### Example A1


a) The preparation of intermediate 1



nBuLi (0.012 mol) was added dropwise at  $-70^{\circ}\text{C}$  to a mixture of *N*'-(4-bromo-2,6-dimethylphenyl)-*N,N*-dimethylmethanimidamide (0.0078 mol) in THF (20 ml) under  $\text{N}_2$  flow. The mixture was stirred at  $-30^{\circ}\text{C}$  for 30 minutes, then cooled to  $-70^{\circ}\text{C}$ . A mixture of DMF (0.078 mol) in THF (30 ml) was added dropwise. The mixture was stirred at

$-70^{\circ}\text{C}$  for 2 hours, then brought to  $0^{\circ}\text{C}$ , poured out into  $\text{H}_2\text{O}$  and extracted with ethyl acetate. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. Yield: 1.8 g of intermediate 1.

b) The preparation of intermediate 2




A mixture of diethyl (cyanomethyl)phosphonate (0.0037 mol) in THF (10 ml) was cooled to  $5^{\circ}\text{C}$  under  $\text{N}_2$  flow. Potassium tert.-butoxide (0.0037 mol) was added portionwise. The mixture was stirred at  $5^{\circ}\text{C}$  for 30 minutes, then stirred at room temperature for 30 minutes. A mixture of intermediate 1 (0.0024 mol) in THF (10 ml) was added. The mixture was stirred at room temperature for 1 hour, then poured out


-54-

into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. Yield : 0.82 g (100%) of intermediate 2.

c) The preparation of intermediate 3 and intermediate 22



Intermediate 3



Intermediate 22

A mixture of intermediate 2 (0.059 mol) and ZnCl<sub>2</sub> (0.299 mol) in ethanol (150 ml) was stirred and refluxed for 24 hours, then poured out into K<sub>2</sub>CO<sub>3</sub> solution (10 % in water) and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (9 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.8 g (6%) of intermediate 22. The filtrate was concentrated and recrystallized from DIPE to obtain 6 g of intermediate 3.

Alternatively, intermediate 3 was also prepared as follows:

To a solution of 159 g of 4-iodo-2,6-dimethyl-benzenamine was added 63.8 g of sodium acetate. The reaction mixture was kept under nitrogen atmosphere. 7 g of moistened palladium on charcoal (Pd/C 10 %) and 64.4 ml of acrylonitrile was added. The reaction mixture was heated to 130°C and stirred overnight. After cooling to room temperature, 0.5 l of toluene and 0.5 l of *N,N*-dimethylacetamide was added. The reaction mixture was filtered over Dicalite and the filter was washed with 0.5 l of toluene. Water (6 l) was added to the mixture which was stirred for 30 minutes. The layers were separated. To the aqueous layer, 1 l of toluene was added and the mixture was stirred for 30 minutes. The layers were separated again. The separated organic layers were collected and the solvent was evaporated, yielding 123 g of intermediate 3.

Intermediate 3 was converted into its hydrochloric acid salt as follows :


To a mixture of 123 g of intermediate 3 in 630 ml of ethanol was added 1,25 l of diisopropyl ether. The reaction mixture was kept under nitrogen atmosphere. The mixture was heated to 60°C and stirred for 30 minutes. 120 ml of a 6 N solution of hydrochloric acid in 2-propanol was added and the mixture was stirred for 30 minutes. After cooling to room temperature, the reaction mixture was filtered and the residue was washed with 100 ml of 2-propanol. The resulting residue was dried under reduced pressure at 50°C. Yield : 103 g (77 %) of the hydrochloric acid salt (1:1) of intermediate 3.



-55-

Intermediate 3 (E) was prepared as follows :


x) Preparation of intermediate 3a (E)



Intermediate 3a (E)

In 10 ml acetonitrile, dry, was dissolved 2.00 g (10.0 mol) of 4-bromo-2,6-dimethylaniline, 1.07 g (1.5 eq) of acrylamide, 224 mg (0.1 eq) of Pd(OAc)<sub>2</sub>, 609 mg (0.2 eq) of tris(2-methylphenyl)phosphine and 1.52 g of *N,N*-diethylethanamine. The mixture was purged with N<sub>2</sub> for 20 minutes and stirred overnight at 70 °C. The mixture was diluted with 150 ml of methylene chloride, washed with saturated aqueous NaHCO<sub>3</sub> solution, dried (sat. NaCl, Na<sub>2</sub>SO<sub>4</sub>) and filtered. The solvent was evaporated and the residue was stirred in diisopropyl ether followed by filtration. Yield : 1.51 g (79.5 %) of intermediate 3a (E).


y) Preparation of intermediate 3 (E)



Intermediate 3 (E)

POCl<sub>3</sub> (3 ml) was cooled to 0°C and 500 mg (2.63 mmol) of intermediate 3a (E) was added. After 30 minutes, the cooling bath was removed and the mixture was stirred overnight at 20 °C. The mixture was added dropwise to 150 ml of diisopropyl ether while stirring vigorously. The precipitate was filtered and washed with diisopropyl ether. The residue was added to 100 ml ethyl acetate/100 ml of saturated aqueous NaHCO<sub>3</sub> solution and stirred. The ethyl acetate layer was separated, dried (sat. NaCl, Na<sub>2</sub>SO<sub>4</sub>) and filtered. The solvent was evaporated. Yield : 380 mg (84 %) of intermediate 3 (E).

d) The preparation of intermediate 4




-56-

A mixture of 4-bromo-2,6-dimethylbenzenamine (0.024 mol) in  $\text{H}_2\text{SO}_4$  (30 ml) was stirred at  $-5^\circ\text{C}$ .  $\text{KNO}_3$  (0.024 mol) was added slowly. The mixture was stirred at  $-5^\circ\text{C}$  for 30 minutes, poured out into  $\text{H}_2\text{O}$  and extracted with ethyl acetate. The organic layer was washed with  $\text{H}_2\text{O}$ , separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was  
5 evaporated. The residue (0.058 g, 95%) was purified by column chromatography over silica gel (eluent: cyclohexane/ethyl acetate; 70/30; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 4.1 g of intermediate 4.

#### Example A1A

The preparation of intermediate 28




1-chloro-pyrrolidine-2,5-dione (0.032 mol) was added at  $60^\circ\text{C}$  to a mixture of 4-amino-  
10 3-methyl-benzoic acid ethyl ester [CAS 40800-65-5] (0.029 mol) in  $\text{CH}_3\text{CN}$  (50 ml). The mixture was stirred and refluxed slowly.  $\text{K}_2\text{CO}_3$  10% was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was evaporated. The residue (6.6 g) was purified by column chromatography over silica gel (eluent: cyclohexane/ $\text{EtOAc}$  85/15; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield:  
15 5.2 g of intermediate 28 (84%).

#### Example A2

A mixture of 4-[(1,4-dihydro-4-oxo-2-pyrimidinyl)amino]benzonitrile (0.12 mol) in  $\text{POCl}_3$  (90 ml) was stirred and refluxed under Argon for 20 minutes. The reaction mixture was slowly poured onto 750 ml ice/water, and the solid was separated by  
20 filtration. The solid was suspended in 500 ml water, and the pH of the suspension was adjusted to neutral by adding a 20%  $\text{NaOH}$  solution. The solid was again separated by filtration, suspended in 200 ml 2-propanone, and 1000 ml  $\text{CH}_2\text{Cl}_2$  was added. The mixture was heated until all solid had dissolved. After cooling to room temperature, the aqueous layer was separated, and the organic layer was dried. During removal of  
25 the drying agent by filtration, a white solid formed in the filtrate. Further cooling of the filtrate in the freezer, followed by filtration, yielded 21.38 g (77.2%) of [4-[(4-chloro-2-pyrimidinyl)amino]benzonitrile (interm. 5).

#### Example A3


a) The preparation of intermediate 6



-57-


nBuLi (0.024 mol) was added dropwise at  $-70^{\circ}\text{C}$  to a mixture of *N'*-(4-bromo-2,6-dimethylphenyl)-*N,N*-dimethylmethanimidamide (0.0157 mol) in THF (50 ml) under  $\text{N}_2$  flow. The mixture was stirred at  $-30^{\circ}\text{C}$  for 30 minutes, then cooled to  $-70^{\circ}\text{C}$ . A solution of 2-methylpropanal (0.055 mol) in THF (50 ml) was added. The mixture was stirred at  $-70^{\circ}\text{C}$  for 2 hours, then brought to  $0^{\circ}\text{C}$ , poured out into  $\text{H}_2\text{O}$  and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (6.7 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  95/5/0.5; 15-40 $\mu\text{m}$ ). Two fractions were collected and the solvent was evaporated. Fraction 1 : yield : 1.5 g of intermediate 6 (38%).

b) The preparation of intermediate 7




Tris[2-(2-methoxyethoxy)ethyl]amine (0.0193 mol) was added at room temperature to a solution of intermediate 6 (0.0048 mol) in  $\text{CH}_2\text{Cl}_2$  (20 ml).  $\text{KMnO}_4$  (0.0193 mol) was added portionwise. The mixture was stirred at room temperature overnight, then filtered over celite and washed with  $\text{CH}_2\text{Cl}_2$ . The organic layer was washed with  $\text{K}_2\text{CO}_3$  10%, separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. Yield: 1.2 g (100%) of intermediate 7.

c) The preparation of intermediate 8



A mixture of intermediate 7 (0.0043 mol) and  $\text{ZnCl}_2$  (0.017 mol) in ethanol (20 ml) was stirred and refluxed overnight, poured out into  $\text{H}_2\text{O}$  and extracted with  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. Yield: 0.94 g (82%) of intermediate 8.


d-1) The preparation of intermediate 9



A mixture of intermediate 8 (0.0049 mol) and intermediate 5 (0.0025 mol) was stirred at  $150^{\circ}\text{C}$  for 2 hours and taken up in  $\text{K}_2\text{CO}_3$  10%/ $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (1.3 g) was crystallized from DIPE. The precipitate was filtered off and dried. The mother layer was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  98.5/1.5; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.21 g of intermediate 9.


-58-

d-2) The preparation of intermediate 29




A mixture of intermediate 28 (0.023 mol) and intermediate 5 (prepared according to A2) (0.025 mol) in HCl 3N (10 ml) was stirred at 105°C then brought to room temperature and filtered. The precipitate was washed with DIPE and dried. Yield: 8.4 g of intermediate 29 (96%)

d-3) The preparation of intermediate 30




- 5 A mixture of 4-amino-3-chloro benzoic acid ethyl ester [CAS 82765-44-4] (0.02 mol) and intermediate 5 (prepared according to A2) (0.0243 mol) in 1-methyl-pyrrolidin-2-one (40 ml) was stirred at 180°C for 2 hours, then poured out into H<sub>2</sub>O and extracted three times with EtOAc (80 ml). The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (10 g)
- 10 was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub> 100; 15-30 μm). Two fractions were collected and the solvent was evaporated. Yield: 1.7 g F1 and 1 g F2. F2 was taken up in diethyl ether. The precipitate was filtered off and dried. Yield: 0.95 g of intermediate 30 (12%).

e-1) The preparation of intermediate 17




- 15 NaBH<sub>4</sub> (0.0001 mol) was added portionwise at 5°C to a mixture of intermediate 9 (0.0001 mol) in ethanol (7 ml) under N<sub>2</sub> flow. The mixture was stirred at 5°C for 1 hour, poured out on ice and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.1 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.044 g of intermediate 17.

e-2) The preparation of intermediate 32



- 20 BuLi 1.6 M (0.009 mol) was added at -78°C to a mixture of




(intermediate 31) (prepared according to A4a) (0.0029 mol) in


-59-

THF (25 ml) under N<sub>2</sub> flow. The mixture was stirred at -78°C for 10 minutes, then brought to room temperature and stirred for 3 hours. H<sub>2</sub>O was added. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (1.28 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 98/2/0.1; 15-40μm). Three fractions were collected and the solvent was evaporated. Yield: 0.189 g of fraction 1, 0.14 g of fraction 2 and 0.5 g of fraction 3 (48%). Fraction 3 was purified by column chromatography over kromasil (eluent: CH<sub>2</sub>Cl<sub>2</sub>/EtOAc 80/20; 10μm). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.25 g F1 (24%) and 0.1 g of F2. F1 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.21 g of intermediate 32 (20%).

e-3) The preparation of intermediate 34




A solution of methyllmagnesium iodide (1.0M solution in diethylether) (0.6ml)

was added to a solution of  intermediate 33 (prepared according to A5.a) (0.0006 mol) in THF (3 ml). The mixture was stirred for 2 hours. H<sub>2</sub>O was added. The mixture was filtered over celite. H<sub>2</sub>O was added. The mixture was extracted with EtOAc. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.05 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 96/4; 15-40μm). The pure fractions were collected and the solvent was evaporated. Yield: 0.015 g of intermediate 34 (7.2%).

#### Example A4

a) The preparation of intermediate 10




A mixture of ethyl 3,5-dimethyl-4-hydroxy benzoate (0.0025 mol) in 1,4-dioxane (2.5 ml) was stirred at room temperature under N<sub>2</sub> flow. Sodium hydride (0.0033 mol) was added. The mixture was stirred for 2 minutes. Intermediate 5 (0.0028 mol) was added. The mixture was stirred for 10 minutes. 1-methyl-2-pyrrolidinone (2.5 ml) was added. The mixture was stirred at 150°C for 12 hours, poured out into H<sub>2</sub>O and extracted with

-60-


$\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (1.7 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  92/8; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.7 g of intermediate 10 (70%).

b-1) The preparation of intermediate 11




- 5 A solution of intermediate 10 (0.0005 mol) in THF (5 ml) was added dropwise at 0°C to a suspension of  $\text{LiAlH}_4$  (0.001 mol) in THF (5 ml) under  $\text{N}_2$  flow. The mixture was stirred at 0°C for 1 hour and poured out into  $\text{H}_2\text{O}$  (0.5 ml).  $\text{CH}_2\text{Cl}_2$  was added. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2$  100 to  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1; 5 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. The residue (0.1 g) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.043 g of intermediate 11 (24%).

b-2) The preparation of intermediate 37



- 15  $\text{LiAlH}_4$  (0.0196 mol, 0.75 g) was added portionwise at 5°C to a mixture of intermediate 29 (prepared according to A3d-2) (0.0098 mol) in THF (100 ml) under  $\text{N}_2$  flow. The mixture was stirred at room temperature overnight, poured out into EtOAc, then into  $\text{H}_2\text{O}$  and filtered over celite. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. Yield : 3.4 g. This fraction was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  97/3/0.1; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield 1 g (27%).
- 20 This fraction was crystallized from DIPE/ $\text{CH}_3\text{CN}$ . The precipitate was filtered off and dried. Yield: 0.03 g of intermediate 37.


c) The preparation of intermediate 12



-61-

A mixture of intermediate 11 (0.0043 mol) in  $\text{CH}_2\text{Cl}_2$  (50 ml) was stirred at  $0^\circ\text{C}$ .  $\text{SOCl}_2$  (0.0206 mol) was added dropwise. The mixture was poured out into ice water/ $\text{K}_2\text{CO}_3$ . The mixture was stirred at room temperature for 5 minutes. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. Yield: 1.5 g of intermediate 12 (98%).


d) The preparation of intermediate 55



Jones's reagent (0.0084 mol) was added to a mixture of intermediate 19 (see Table 1) (prepared according to A4b-1) (0.0028 mol) in acetone (50 ml). The mixture was stirred at room temperature for 2 hours then poured out into  $\text{H}_2\text{O}$  and basified with  $\text{NaHCO}_3$ . The precipitate was filtered off and dried. Yield : 1.39 g. The residue (0.1 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  85/15/1 then  $\text{CH}_3\text{OH}$  100). The pure fraction was crystallized from isopropanol/DIPE. Yield : 0.071 g of intermediate 55.


#### Example A5

a) The preparation of intermediate 13



A mixture of intermediate 19 (see Table 1) (prepared according to A4.b-1) (0.0037 mol) and  $\text{MnO}_2$  (0.0185 mol) in  $\text{CH}_2\text{Cl}_2$  (100 ml) was stirred at room temperature overnight, then filtered over celite. The filtrate was evaporated. Yield: 1.3 g of intermediate 13.

b) The preparation of intermediate 21




A mixture of intermediate 13 (prepared according to A5.a) (0.0029 mol) and  $\text{H}_2\text{N}-\text{NH}_2$ ,  $\text{H}_2\text{O}$  (0.0058 mol) in EtOH (10 ml) was stirred at room temperature overnight. The solvent was evaporated till dryness. Yield: 0.53 g of intermediate 21.

Example A6

-62-


The preparation of intermediate 14



Hydrazine (0.0077 mol) was added to a mixture of according to A3.d-1) (0.0005 mol) in EtOH (10 ml). The mixture was stirred and refluxed overnight. Hydrazine (0.028 mol) was added. The mixture was stirred and refluxed overnight. Yield: 0.28 g of intermediate 14.


#### 5 Example A7

a) The preparation of intermediate 23



A mixture of intermediate 35 (0.0056 mol) in HCl 3N (60 ml) and *i*PrOH (15 ml) was stirred and refluxed overnight. The precipitate was filtered, washed with H<sub>2</sub>O, taken up in DIPE and dried. Yield: 2.3 g of intermediate 23 (100%).

b) The preparation of intermediate 56



10 A mixture of intermediate 10 (prepared according to A4.a) (0.0012 mol) in HCl 3N (26 ml) and *i*PrOH (4 ml) was stirred and refluxed for 12 hours. The solvent was evaporated till dryness. The residue was taken up in (CH<sub>3</sub>)<sub>2</sub>CO. The solvent was evaporated. The residue was taken up in diethyl ether. The precipitate was filtered off and dried. Yield: 0.4 g (78.5%). This fraction was stirred at 60°C for 20 minutes.




-63-

Yield: 0.19 g. This fraction was crystallized from H<sub>2</sub>O/2-propanone. The precipitate was filtered off and dried. Yield: 0.12 g of intermediate 56 (26%).


#### Example A8

a) The preparation of intermediate 24



A mixture of intermediate 31 (prepared according to A4.a) (0.0005 mol) and (triphenylphosphoranylidene)acetic acid ethyl ester [CAS 1099-45-2] (0.0006 mol in THF (5 ml) was stirred at 80°C for 48 hours, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (0.4 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.08 g (33%). This fraction was crystallized from DIPE/CH<sub>3</sub>CN. The precipitate was filtered off and dried. Yield: intermediate 24 (33%).


b) The preparation of intermediate 25




Piperidine (0.0011 mol) was added at room temperature for 30 minutes. Intermediate 31 (prepared according to A4.a) (0.0005 mol) was added. The mixture was stirred at room temperature for 1 hour, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The precipitate was filtered off and dried. The residue (0.2 g) was crystallized from CH<sub>3</sub>CN/DIPE. The precipitate was filtered off and dried. Yield: 0.048 g of intermediate 25 (19%) (mp. 222°C).

#### Example A9

The preparation of intermediate 26




-64-



- A mixture of (prepared according to A3.d-1) (0.0011 mol) and Pd/C (0.2 g) in methanol (30 ml) was hydrogenated at room temperature for 2 hours under one bar pressure, then filtered over celite. Celite was washed with CH<sub>3</sub>OH. The filtrate was evaporated till dryness. The residue (0.3 g) was crystallized from 2-propanone/CH<sub>3</sub>OH/diethyl ether. The precipitate was filtered off and dried. Yield: 0.07 g of fraction 1. Fraction 1 was purified by column chromatography over kromasyl (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99.5/0.5; 5 $\mu$ m). Three fractions 9F1, F2, F3) were collected and the solvent was evaporated. Yield: 0.0516 g F1, 0.1g F2 and 0.15 g F3. F1 was taken up in diethyl ether. The precipitate was filtered off and dried. Yield: 0.028 g of intermediate 26 (8%) (mp. 272°C).

#### Example A10


The preparation of intermediate 27



- A mixture of (prepared according to A4.c) (0.0005 mol) and triphenylphosphine (0.0005 mol) in CH<sub>3</sub>CN (10 ml) was stirred and refluxed for a week end. The solvent was evaporated till dryness. The residue was taken up in diethyl ether. The precipitate was filtered off and dried. Yield: 0.34 g of intermediate 27 (94%).

#### Example A11

The preparation of intermediate 58




- A mixture of 4-bromo-2,6-dimethylbenzenamine (0.013 mol) and intermediate 5 (0.013 mol) was stirred at 150°C for 1 hour. The mixture was poured into K<sub>2</sub>CO<sub>3</sub> 10% aqueous solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>/ MeOH (95/5). The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue was crystallized from diisopropyl ether. The precipitate was filtered off and dried. Yield : 2.3g (45%). The mother layer was purified by column chromatography over silica gel (eluent:

-65-

$\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}-\text{NH}_4\text{OH}$  98.5/1.5; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.90 g (17%). The global yield of intermediate 5 was: 3.2 g (62%).


- 5 Intermediate 59 was prepared analogously.



Intermediate 59

Table 1 and 2 list intermediates which intervene in the preparation of compounds of the present invention.

Table 1




5

Interm. No.	Ex. No.	X <sub>1</sub>	R <sup>3</sup>	R <sup>4a</sup>	R <sup>4b</sup>	Physical data
11	A4b-1	O	-CH <sub>2</sub> -OH	CH <sub>3</sub>	CH <sub>3</sub>	
12	A4c	O	-CH <sub>2</sub> -Cl	CH <sub>3</sub>	CH <sub>3</sub>	
16	A3e	NH	-CH(OH)-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	
17	A3e	NH	-CH(OH)-CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	
18	A3e	NH	-CH(OH)-CH <sub>2</sub> -CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	
19	A4b-1	NH	-CH <sub>2</sub> -OH	CH <sub>3</sub>	CH <sub>3</sub>	
15	A4c	NH	-CH <sub>2</sub> -Cl	CH <sub>3</sub>	CH <sub>3</sub>	
24	A8a	O	-CH=CH-C(=O)-O-C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	CH <sub>3</sub>	mp. 180°C; (E)
25	A8b	O		CH <sub>3</sub>	CH <sub>3</sub>	mp. 222°C; (A)
35	A3d-1	NH	-CH=CH-C(=O)-O-C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	CH <sub>3</sub>	mp. 200°C; (E)
23	A7a	NH	-CH=CH-COOH	CH <sub>3</sub>	CH <sub>3</sub>	
34	A3e-3	NH	-CH(OH)-CH <sub>3</sub>	CH <sub>3</sub>	H	mp. 182°C
36	A4b-1	NH	-CH <sub>2</sub> -OH	CH <sub>3</sub>	H	mp. 210°C
37	A4b-2	NH	-CH <sub>2</sub> -OH	Cl	CH <sub>3</sub>	
38	A4b-1	NH	-CH <sub>2</sub> -OH	Cl	H	mp. 226°C
39	A3e-1	O	-CH(OH)-CH <sub>3</sub>	CH <sub>3</sub>	H	mp. 160°C
40	A4b-1	S	-CH <sub>2</sub> -OH	CH <sub>3</sub>	CH <sub>3</sub>	mp. 173°C
41	A4b-1	NH	-CH <sub>2</sub> -OH	Br	H	mp. 234°C
32	A3e-2	O	-CH(OH)-CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	mp. 193°C
42	A4b-1	NH	-CH <sub>2</sub> -OH	Br	CH <sub>3</sub>	mp. 250°C
43	A4b-1	NH	-CH <sub>2</sub> -OH	OH	H	mp. 124°C
44	A4b-1	NH	-CH <sub>2</sub> -OH	H	H	mp. 215°C
45	A4b-1	NH	-CH <sub>2</sub> -OH	O-CH <sub>3</sub>	H	
46	A4b-1	NH	-CH <sub>2</sub> -OH	CF <sub>3</sub>	H	mp. 194°C
47	A4c	NH	-CH <sub>2</sub> -Cl	Cl	CH <sub>3</sub>	
48	A4c	NH	-CH <sub>2</sub> -Cl	Cl	H	

-67-

Interm. No.	Ex. No.	X <sub>1</sub>	R <sup>3</sup>	R <sup>4a</sup>	R <sup>4b</sup>	Physical data
49	A3e-1	O	-CH <sub>2</sub> -OH	CH <sub>3</sub>	H	
50	A4c	O	-CH <sub>2</sub> -Cl	CH <sub>3</sub>	H	
51	A4b-1	NH	-CH <sub>2</sub> -OH	C(CH <sub>3</sub> ) <sub>3</sub>	H	
52	A4c	NH	-CH <sub>2</sub> -Cl	CH <sub>3</sub>	H	
53	A4b-1	NH	-CH <sub>2</sub> -OH	2-furanyl	CH <sub>3</sub>	
54	A4c	NH	-CH <sub>2</sub> -Cl	Br	CH <sub>3</sub>	
57	A7b	O	-CH=CH-COOH	CH <sub>3</sub>	CH <sub>3</sub>	

Table 2




Interm. No.	Ex. No.	X <sub>1</sub>	R <sup>3</sup>	Physical data
20	A3e	NH	-CHOH-CH <sub>3</sub>	

## 5 B. Preparation of the final compounds

### Example B1

The preparation of compound 1



A mixture of intermediate 3 (0.034 mol) and intermediate 5 (0.0174 mol) was stirred at 150°C for 1 hour and taken up in K<sub>2</sub>CO<sub>3</sub> 10%/CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (10 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/ethyl acetate 80/20; 15-40μm). Fraction 1 was crystallized from iPrOH. The precipitate was filtered off and dried. Yield: 1.3 g of 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]amino]-2-pyrimidinyl]amino]benzonitrile (E) (compound 1) (20%).

### 15 Example B1A

Compound 1 was also prepared as follows :

A mixture of 93.9 g (0.45 mol) of the hydrochloric acid salt of intermediate 3, prepared according to Example A1c), and 109 g (0.4725 mol) of intermediate 5 in 1.8 l of

-68-

acetonitrile was prepared under nitrogen atmosphere. The mixture was stirred and refluxed for 69 hours, then allowed to cool to 55 °C. The mixture was filtered and the residue was washed with 200 ml of acetonitrile, followed by drying under reduced pressure at 50°C overnight. 144,6 g (0.3666 mol) of the obtained solid was brought in 1 l of K<sub>2</sub>CO<sub>3</sub> 10% aqueous solution. The mixture was stirred at room temperature followed by filtration. The obtained residue was washed twice with water followed by drying at 50°C under reduced pressure. The residue was brought in 6.55 l isopropanol and the mixture was refluxed, then stirred overnight and filtered at room temperature. The residue was dried at 50°C under reduced pressure. Yield : 113.2 g (68.6 %) of 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]amino]-2-pyrimidinyl]amino]benzonitrile (E) (compound 1).

#### Example B1B

Alternatively, compound 1 was also prepared as follows :


- 15 a) A mixture of intermediate 58 (0.00021 mol), prepared according to Example A11, acrylonitrile (CH<sub>2</sub>=CH-CN) (0.00213 mol), Pd(OAc)<sub>2</sub> (0.000043 mol), *N,N*-diethylethanamine (0.000043 mol) and tris(2-methylphenyl)phosphine (0.00021 mol) in CH<sub>3</sub>CN (7 ml) was stirred in a sealed vessel at 150°C overnight. H<sub>2</sub>O was added. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>),  
20 filtered and the solvent was evaporated. The residue (0.15 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/ethyl acetate 80/20; 15-40 µm). Fraction 1 was collected and the solvent was evaporated, yielding 0.045g of 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]amino]-2-pyrimidinyl]amino]benzonitrile (E/Z=80/20). The solid was crystallized from diethylether. Yield: 0.035g of 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]amino]-2-pyrimidinyl]amino]benzonitrile (E) (compound 1) (55%).  
25 b) 4,41g (10 mmol) of intermediate 59 and 15 ml of *N,N*-dimethylacetamide were brought in a 100 ml flask under nitrogen. To this mixture were added 0,98g of sodium acetate (12 mmol), 107 mg (0,1 mmol Pd) of Pd/C 10% (wet) and 1 ml (15 mmol) of acrylonitrile. The mixture was heated at 140°C and the evolution of the reaction was followed by liquid chromatography. The reaction yielded 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]amino]-2-pyrimidinyl]amino]benzonitrile (E/Z=80/20) which can be converted to 4-[[4-[[4-(2-cyanoethenyl)-2,6-dimethylphenyl]amino]-2-pyrimidinyl]amino]benzonitrile (E) as described above in Example B1Ba).


35

#### Example B2


-69-


## a) The preparation of compound 2




A mixture of  (prepared according to A3.d-1) (0.0002 mol), 2-benzofuranylboronic acid (0.0005 mol),  $\text{Pd(PPh}_3)_4$  (0.00002 mol) and  $\text{Na}_2\text{CO}_3$  (0.0007 mol) in DME (3 ml) was stirred and refluxed in a sealed tube for 3 hours.  $\text{H}_2\text{O}$  was added. The mixture was extracted with ethyl acetate. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (0.126 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  98/2; 15-40  $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.011 g of compound 2 (10%).



## b) The preparation of compound 3



A mixture of  (prepared according to A3.d-1) (0.0002 mol), tributyl-2-furanylstannane (0.0005 mol) and  $\text{Pd(PPh}_3)_4$  (0.00001 mol) in dioxane (5 ml) was stirred at 80°C. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  98/2; 15-40  $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. The residue (0.025 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.021 g of compound 3 (22%).

## c) The preparation of compound 104




A mixture of  (prepared according to A3.d) (0.005 mol),  [CAS 73183-34-3] (0.0055 mol),  $\text{Pd(PPh}_3)_4$  (0.29 g) and  $\text{K}_2\text{CO}_3$  (2.8 g, 0.02 mol) in toluene (100 ml) and ethanol/water (5 to 10 ml) was stirred and refluxed for a weekend. 5-Bromo-furan-2-carbaldehyde (0.0055 mol) and  $\text{K}_2\text{CO}_3$  (1.4 g, 0.01

-70-

mol) were added. The mixture was stirred and refluxed overnight. The mixture (2.25 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 to 99/1; 15-40 mm). The pure fractions were collected and the solvent was evaporated. Yield: 0.135 g of compound 104 (6%).

5 Example B3



The preparation of compound 4



A mixture of intermediate 15 (see Table 1) (prepared according to A4.c) (0.0005 mol) and NaCN (0.0011 mol) in DMF (5 ml) was stirred at 80°C overnight, poured out into  $\text{H}_2\text{O}$  and extracted with ethyl acetate. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue (0.15 g) was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1; 10  $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. The residue (0.024 g) was purified by column chromatography over hypersil (eluent: acetonitrile/ $\text{H}_2\text{O}$  52/48; 8  $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.02 g of compound 4 (10%).

15 Example B4

a) The preparation of compound 5




A mixture of (prepared according to A3.d) (0.0006 mol) and thiomorpholine (0.5 g) was stirred at 120°C for 48 hours, taken up in  $\text{CH}_2\text{Cl}_2$  and



-71-

the solvent was evaporated. The residue (0.44 g) was purified by column chromatography over kromasyl (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1;  $10\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.06 g (20%). This fraction was crystallized from diethyl ether/2-propanone. The precipitate was filtered off and dried. Yield: 0.035 g of compound 5.


b) The preparation of compound 6



A mixture of intermediate 15 (see Table 1) (prepared according to A4.c) (0.000137 mol), *N,N,N'*-trimethyl-1,2-ethanediamine (2 equiv, 0.000275 mol) and  $\text{K}_2\text{CO}_3$  (2 equiv, 0.000275 mol) in  $\text{CH}_3\text{CN}$  (q.s.) was stirred at  $80^\circ\text{C}$  for 12 hours.  $\text{H}_2\text{O}$  was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The extract's solvent was evaporated.

10 The residue was purified by chromatography. The product fractions were collected and the solvent was evaporated. Yield: 0.006 g of compound 6 (10.16%).


c) The preparation of compound 7



A mixture of intermediate 15 (see Table 1) (prepared according to A4.c) (0.0005 mol) in 3-hydroxy-propanenitrile (2 ml) was stirred overnight, poured out into  $\text{H}_2\text{O}$  and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  99/1/0.1;  $15\text{--}40\mu\text{m}$ ). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.034 g F1 and 0.514 g F2. F2 was washed with  $\text{HCl}$  3N and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.039 g of compound 7 (18%)

-72-


## d) The preparation of compound 105



A mixture of intermediate 50 (prepared according to A4c) (0.001 mol), KCN (0.0011 mol) and KI (0.00005 mol) in EtOH (15 ml) was stirred and refluxed for 4 hours. The solvent was evaporated till dryness. The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.31 g) was purified by column chromatography over kromasil (eluent: cyclohexane/EtOAc 70/30; 10 $\mu$ m). Three fractions were collected and the solvent was evaporated. Yield: 0.044 g of fraction 1, 0.11 g of fraction 2 and 0.055 g of fraction 3. Fraction 3 was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.046 g of compound 105 (12%) (mp. 140°C).


Example B5

## a) The preparation of compound 8



A mixture of intermediate 9 (0.0001 mol) and hydroxylamine (0.0002 mol) in EtOH (7 ml) was stirred at room temperature for 3 hours, poured out into K<sub>2</sub>CO<sub>3</sub> 10% and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (0.1 g) was crystallized from DIPE/CH<sub>3</sub>CN. The precipitate was filtered off and dried. Yield: 0.026 g of compound 8.

## b) The preparation of compound 9




A mixture of intermediate 9 (0.0002 mol) and O-methylhydroxylamine (0.0003 mol) in EtOH (10 ml) was stirred at room temperature overnight, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (0.13 g) was purified by column chromatography over kromasil (eluent: cyclohexane/iPrOH/NH<sub>4</sub>OH; 5 $\mu$ m). The pure fractions were

-73-

collected and the solvent was evaporated. The residue (0.06 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.036 g of compound 9 (34%).

### Example B6


a) The preparation of compound 1 and 10



Compound 1=(E); compound 10=(Z)

5 A mixture of (cyanomethyl)triphenylphosphonium chloride (0.0022 mol) and potassium tert.-butoxide (0.0022 mol) in THF (7 ml) was stirred at 5°C for 30 minutes under N<sub>2</sub> flow, then stirred at 5°C for 30 minutes. A mixture of intermediate 13 (0.0015 mol) in THF (7 ml) was added. The mixture was stirred for 8 hours in darkness, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (1.4 g) was purified by column chromatography over silica gel (eluent: toluene/iPrOH/NH<sub>4</sub>OH 96/4/0.1; 15-40μm).  
 10 Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.165 g of F1 (E/Z=32/68) (30%) and 0.225 g of F2 (E/Z=90/10) (41%). F2 was crystallized from CH<sub>3</sub>CN/diethyl ether. Yield: 0.036 g of compound 1 (7%). F1 was purified by column chromatography over kromasyl (eluent: toluene/iPrOH 98/2; 5μm). The pure fractions  
 15 were collected and the solvent was evaporated. Yield: 0.029 g of compound 10 (5%).


b) The preparation of compound 11 (Z) and compound 103 (E)



Potassium tert-terbutoxide (0.0196 mol) was added portionwise at 5°C to a mixture of (1-cyanoethyl)-phosphonic acid diethyl ester (0.0196 mol) in THF (25 ml) under N<sub>2</sub> flow. The mixture was stirred at 5°C for 30 minutes, then at room temperature for 30 minutes. A solution of intermediate 13 (0.0130 mol) in THF (25 ml) was added. The  
 20 mixture was stirred at room temperature overnight, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (5.8 g) was purified by column chromatography over silica gel (eluent: toluene/iPrOH/NH<sub>4</sub>OH 92/8/0.5; 15-40μm). Four fractions (F1, F2, F3, F4) were collected and the solvent was evaporated. Yield: 0.21 g of F1 (mixture  
 25 Z/E=90/10), 0.836 g of F2 (mixture Z/E=57/43), 0.9 g of F3 and 0.87 g of F4. F3 was crystallized from DIPE/iPrOH to give 0.7 g of compound 11 (14%). F4 was crystallized from DIPE/ iPrOH to give 0.67 g of compound 103 (13%).


-74-

## c) The preparation of compound 12 and 13



compound 12 = (E)  
compound 13 = (Z)


Potassium tert.-butoxide (0.0008 mol) was added portionwise at 5°C to a mixture of (cyanomethyl)phosphonic acid diethyl ester (0.0005 mol) in THF (20 ml) under N<sub>2</sub> flow. The mixture was stirred at room temperature for 30 minutes. A solution of



5 ml) was added dropwise. The mixture was stirred at room temperature for 4 hours, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. Yield: 0.3 g. This fraction was purified by column chromatography over kromasil (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 5μm). The pure fractions were collected and the solvent was evaporated. Yield: 0.21 g. This


10 fraction was purified by column chromatography over kromasil (eluent: cyclohexane/ethyl acetate 50/50; 10μm). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.04 g of F1 and 0.047 g F2. F1 was dried at 70°C for 2 hours. Yield: 0.038 g of compound 13 (18%). F2 was dried at 70°C for 2 hours. Yield: 0.041 g of compound 12 (20%).

## d) The preparation of compound 14




-75-

Potassium tert.-butoxide (0.0013 mol) was added at 5°C to a mixture of (cyanomethyl)phosphonic acid diethyl ester (0.0013 mol) in THF (10 ml) under N<sub>2</sub> flow. The mixture was stirred at 5°C for 30 minutes. A mixture of



- (prepared according to A3.d-1) (0.0009 mol) in THF (10 ml) was added. The mixture was stirred at room temperature for 4 hours, poured out into H<sub>2</sub>O and extracted with ethyl acetate. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.17 g) was purified by column chromatography over kromasil (eluent: CH<sub>2</sub>Cl<sub>2</sub> 100 to CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 5μm). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.054 g F1 and 0.05 g F2. F1 was crystallized from DIPE/CH<sub>3</sub>CN. The precipitate was filtered off and dried. Yield: 0.046 g of compound 14 (12%).


e) The preparation of compound 15



- 4-Fluorobenzeneacetonitrile (1.2 equiv, 0.000175 ml) was added to a mixture of intermediate 13 (0.000146 mol) in CH<sub>3</sub>OH (1 ml). NaOCH<sub>3</sub>/CH<sub>3</sub>OH (1.2 equiv, 0.000175 mol) was added at room temperature. The mixture was stirred at 60 °C for 2 hours, then poured out into ice-water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The solvent was evaporated. The residue was purified by chromatography. The product fractions were collected and the solvent was evaporated. Yield: 0.009 g of compound 15 (13.42%).

-76-

## f) The preparation of compound 106




A mixture of intermediate 13 (prepared according to A5.a) (0.0005 mol) and piperidine (0.0005 mol) in ethanol (5 ml) was stirred at room temperature for 30 minutes. 4,4-dimethyl-3-oxo-pentanenitrile (0.0011 mol) was added. The mixture was stirred at room temperature overnight, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.3 g) was purified by column chromatography over kromasil (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 10 $\mu$ m). The pure fractions were collected and the solvent was evaporated. The residue (0.2 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.141 g of compound 106 (54%) (mp. 193°C).

10

Example B7

The preparation of compound 16




15

A mixture of intermediate 14 (0.00005 mol) and carbonothioic dichloride (0.001 mol) in dioxane (10 ml) was stirred at room temperature. H<sub>2</sub>O was added. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. This fraction was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 90/10/0.1; 15-40 $\mu$ m). The pure fractions were collected and the solvent was evaporated. Yield: 0.027 g of compound 16 (95.6%).


-77-

Example B8

The preparation of compound 17



The mixture of NaOCH<sub>3</sub> (0.001 mol) and 2-(dimethylamino)-*N*-hydroxyethanimidamide (0.001 mol) in EtOH (10 ml) was stirred at room temperature for 30




minutes.

(prepared according to A3.d-1) (0.0005 mol) was added.

- 5 The mixture was stirred and refluxed overnight. H<sub>2</sub>O was added. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 95/5/0.1; 15-40 μm). The pure fractions were collected and the solvent was evaporated. Yield: 0.07 g of compound 17 (31%).

Example B9

The preparation of compound 18




- 10 nBuLi (0.0038 mol) was added dropwise at -70°C to a mixture of iPr<sub>2</sub>NH (0.0038 mol) in THF (5 ml) under N<sub>2</sub> flow. The mixture was brought to -20°C, stirred for 30 minutes and cooled again to -70°C. A solution of CH<sub>3</sub>CN (0.0038 mol) in THF (6 ml) was added dropwise. The mixture was brought to -20°C, stirred for 1 hour, cooled again to -70°C. A mixture of intermediate 13 (0.0009 mol) in THF (1 ml) was added. The
- 15 the mixture was stirred for 2 hours, poured out on ice at -30°C and extracted with ethyl acetate. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was

-78-

evaporated. The residue (0.433 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  98/2; 35-70 $\mu\text{m}$ ). Two fractions were collected and the solvent was evaporated. Yield: 0.056 g F1 and 0.23 g F2 (78%). F1 was crystallized from DIPE/ $\text{CH}_3\text{CN}$ . The precipitate was filtered off and dried. Yield: 0.036 g of compound 18.


Example B9A

a) The preparation of compound 107




nBuLi[1.6] (0.0026 mol) was added dropwise at  $-70^\circ\text{C}$  to a mixture of intermediate 13 (prepared according to A5.a) (0.0008 mol) in THF (10 ml) under  $\text{N}_2$  flow. The mixture was stirred at  $-70^\circ\text{C}$  for 30 minutes. A solution of (chloromethyl)triphenylphosphonium chloride (0.0026 mol) in THF (5 ml) was added dropwise. The mixture was stirred at room temperature overnight, poured out into  $\text{H}_2\text{O}$  and extracted with EtOAc. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (0.7 g) was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1; 10 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. The residue (0.155 g) was purified by column chromatography over C18 (eluent:  $\text{CH}_3\text{CN}/\text{NH}_4\text{Ac}$  0.5% 60/40). The pure fractions were collected and the solvent was evaporated. The residue (0.051 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.029 g of compound 107 (9%). (mp.  $250^\circ\text{C}$ )

b) The preparation of compound 108 and 109



(Z)

compound 108



(E)

compound 109


nBuLi[1.6] (0.00261 mol) was added dropwise at  $-70^\circ\text{C}$  to a mixture of (chloromethyl)triphenylphosphonium chloride (0.00261 mol) in THF (10 ml) under  $\text{N}_2$



-79-

flow. The mixture was stirred for 30 minutes. A solution of intermediate 31 (prepared according to A4.a) (0.00087 mol) in THF (5 ml) was added dropwise. The mixture was stirred at room temperature overnight, then poured out into H<sub>2</sub>O and extracted with EtOAc. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (1.1 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 98/2/0.1; 15-40 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.3 g) was purified by column chromatography over hypersil C18 (eluent: CH<sub>3</sub>OH/NH<sub>4</sub>Ac 0.5% 70/30). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.097 g F1 and 0.085 g F2. F1 was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.045 g of compound 108 (14%) (mp. 165°C). F2 was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.049 g of compound 109 (15%) (mp. 200°C).


c) The preparation of compound 110



nBuLi[1.6] (1.1 ml, 0.0017 mol) was added dropwise at -70°C to a mixture of 1,1,1,3,3,3-hexamethyldisilazane (HN(TMS)<sub>2</sub>) (0.0017 mol) in THF (6 ml). The mixture was stirred at -70°C for 30 minutes. Cyanofluoromethyl (0.0017 mol) was added. The mixture was stirred for 30 minutes. Phosphorochloridic acid diethyl ester (0.0017 mol) was added. The mixture was stirred at -70°C for 15 minutes. nBuLi[1.6] (1.1 ml, 0.0017 mol) was added dropwise. The mixture was stirred for 30 minutes. A solution of intermediate 31 (prepared according to A4.a) (0.0008 mol) in THF (4 ml) was added. The mixture was stirred at room temperature overnight, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.5 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/EtOAc 95/5; 15-40 μm). Four fractions (F1, F2, F3, F4) were collected and the solvent was evaporated. Yield: 0.026 g of compound 110 (8%) (mp. 254°C).

-80-


## d) The preparation of compound 111



A solution of  $(\text{CuCl})_2$  (0.00015 mol) in  $\text{NH}_3$  aqueous ( $500\mu\text{l}$ ) was added to a mixture of intermediate 21 (prepared according to A5.b) (0.0014 mol) in DMSO (1 ml). A solution of  $\text{CBr}_4$  (0.0044 mol) in DMSO (1.5 ml) was added at  $0^\circ\text{C}$ . The mixture was stirred at room temperature overnight, poured out on ice and filtered. The organic layer was washed with  $\text{CH}_2\text{Cl}_2$ , dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (2.73 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 to 99/1; 15-40 $\mu\text{m}$ ). Two fractions were collected and the solvent was evaporated. Yield: 0.007 g of fraction 1 and 0.11 g of fraction 2. Fraction 2 was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.075 g of compound 111 (mp.  $223^\circ\text{C}$ ).

Example B9B

## a) The preparation of compound 112




A mixture of intermediate 23 (0.0005 mol), 1-hydroxybenzotriazole (0.0007 mol) and EDCI (0.0007 mol) in  $\text{CH}_2\text{Cl}_2$  (10 ml) and THF (2 ml) was stirred. A solution of  $\text{NH}(\text{CH}_3)_2\cdot\text{HCl}$  (0.0006 mol) and  $\text{Et}_3\text{N}$  (0.0005 mol) was added. The mixture was stirred at room temperature for 12 hours.  $\text{H}_2\text{O}$  was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 to 90/10; 5 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.124 g (58%). This fraction was purified by column

-81-


chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  99/1;  $5\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.045 g of compound 112 (21%) (mp.  $>264^\circ\text{C}$ ).

b) The preparation of compound 113



A mixture of intermediate 57 (prepared according to A7.b) (0.0002 mol),  
 5 1-hydroxybenzotriazole (0.0003 mol) and EDCI (0.0003 mol) in  $\text{CH}_2\text{Cl}_2$  (10 ml) was stirred. *N*-methyl-1-butanamine [CAS 110-68-9] (0.0002 mol) was added. The mixture was stirred at room temperature for 12 hours.  $\text{H}_2\text{O}$  was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. Yield: 0.149 g. This fraction was purified by column  
 10 chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 to 90/10;  $5\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.065 g. This fraction was taken up in DIPE. The precipitate was filtered off and dried. Yield: 0.035 g of compound 113 (30%) (mp.  $212^\circ\text{C}$ ).

c) The preparation of compound 114




A mixture of intermediate 23 (prepared according A7.a) (0.0005 mol),  
 15 1-hydroxybenzotriazole (0.0007 mol) and EDCI (0.0007 mol) in  $\text{CH}_2\text{Cl}_2$  (10 ml) and THF (2 ml) was stirred. 3-(methylamino)propanenitrile (0.0006 mol) was added. The mixture was stirred at room temperature for 12 hours.  $\text{H}_2\text{O}$  was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered,

-82-

and the solvent was evaporated. The residue was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 to 90/10;  $5\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.068 g. This fraction was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.032 g of compound 114 (14%) (mp.  $168^\circ\text{C}$ ).


d) The preparation of compound 115



A mixture of (0.000195 mol) and methylamine (2 equiv, 0.000390 mol) in THF (5 ml) and  $\text{Et}_3\text{N}$  (0.054 ml) was stirred at room temperature. EDCI (2 equiv, 0.000390 mol) and 1-hydroxy-benzotriazole (2 equiv, 0.000390 mol) were added. The reaction mixture was stirred at room temperature for 12 hours and taken up into  $\text{H}_2\text{O}$ . The organic layer was separated, dried, filtered and the solvent evaporated. The product was isolated and purified by column chromatography. Yield: 0.026 g of compound 115 (17.92%).

#### Example B9C

The preparation of compound 116




A mixture of intermediate 13 (prepared according to A5.a) (0.000291 mol) and isonicotinic acid hydrazide (2.5 equiv., 0.000728 mol) in ethanol (1 ml) and  $\text{CH}_2\text{Cl}_2$

-83-

(2 ml) was stirred and refluxed for 12 hours. The solvent was evaporated till dryness. The residue was purified by chromatography. Yield: 0.033 g of compound 116 (24.50%).



### Example B9D

a) The preparation of compound 117



- 5 Sodium cyanoborohydride (0.0024 mol) was added at room temperature to a solution of intermediate 26 (prepared according to A9) (0.0008 mol) in formaldehyde (0.5 ml) and  $\text{CH}_3\text{CN}$  (20 ml) under  $\text{N}_2$  flow. Acetic acid (0.5 ml) was added. The mixture was stirred at room temperature for 2 hours, poured out into  $\text{H}_2\text{O}/\text{K}_2\text{CO}_3$  10% and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue (0.3 g) was purified by column chromatography over hypersol (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  97/3;  $5\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.08 g (28%). This fraction was crystallized from 2-propanone/diethyl ether. The precipitate was filtered off and dried. Yield: 0.012 g of compound 117 (5%) (mp.  $132^\circ\text{C}$ ).

b) The preparation of compound 118



- 15 A mixture of (prepared according to A9) (0.0015 mol) and tetrahydro-2,5-dimethoxyfuran (0.0077 mol) in acetic acid (10 ml) was stirred and refluxed for 1 hour, then poured out into ice water and  $\text{K}_2\text{CO}_3$  and extracted with


-84-

$\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue (1 g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 95/5; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.23 g. This fraction was crystallized from


- 5 DIPE/diethyl ether. The precipitate was filtered off and dried. Yield: 0.075 g. This fraction was crystallized again from DIPE/diethyl ether. The precipitate was filtered off and dried. Yield: 0.027 g of compound 118 (5%).

#### Example B9E

a) The preparation of compound 119




- 10 Tributylphosphine (0.0015 mol) was added to a mixture of but-2-enedinitrile (0.0015 mol) in THF (8 ml). The mixture was stirred and refluxed for 2 hours.



- 15 prepared according to A5.a) (0.0005 mol) was added. The mixture was stirred and refluxed overnight.  $\text{H}_2\text{O}$  was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered and the solvent was evaporated. The residue (0.618 g) was purified by column chromatography over kromasil (eluent:  $\text{CH}_2\text{Cl}_2$  100; 10  $\mu\text{m}$ ). Two fractions were collected and the solvent was evaporated. Yield : 0.03 g of compound 119 (13%).

b) The preparation of compound 120




-85-

Intermediate 13 (prepared according to A5.a) (0.002 mol) was added to a mixture of propanedinitrile (0.004 mol) and piperidine (0.004 mol) in ethanol (10 ml). The mixture was stirred at room temperature for 5 minutes. The solvent was evaporated. The residue was taken up in  $\text{CH}_2\text{Cl}_2$  and purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  98/2; 15-40 $\mu\text{m}$ ). The pure fractions were collected and the solvent was evaporated. Yield: 0.6 g of compound 120.

#### Example B9F


The preparation of compound 122



nBuLi [1.6 M] (0.0016 mol) was added dropwise at  $-78^\circ\text{C}$  to a mixture of intermediate 27 (prepared according to A10) (0.0004 mol) in THF (10 ml) under  $\text{N}_2$  flow. The mixture was stirred at  $-78^\circ\text{C}$  for 1 hour, then brought to room temperature, stirred for 30 minutes and cooled to  $-78^\circ\text{C}$ . A solution of 2-pyridinecarboxaldehyde (0.0004 mol) in THF (10 ml) was added. The mixture was stirred at room temperature for 2 hours, poured out on ice and extracted with EtOAc. The organic layer was separated, dried ( $\text{MgSO}_4$ ), filtered, and the solvent was evaporated. The residue (0.32 g) was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$  98/2/0.1; 10 $\mu\text{m}$ ). Two fractions were collected and the solvent was evaporated. Yield: 0.021 g of compound 122 (10.4%) (mp.  $120^\circ\text{C}$ ).

#### Example B10

The preparation of compound 20




$\text{NaBH}_4$  (0.0015 mol) was added portionwise at  $5^\circ\text{C}$  to a mixture of compound 19 (see table 3) (prepared according to B1) (0.0014 mol) in  $\text{CH}_3\text{OH}$  (15 ml) under  $\text{N}_2$  flow.

-86-

The mixture was stirred at 5°C for 1 hour, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.15 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 10μm). The pure fractions were collected and the solvent was evaporated. The residue (0.068 g, 12%) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.032 g of compound 20.

#### Example B11


The preparation of compound 21



A mixture of compound 2 (see table 3) (0.0002 mol), 3-thienylboronic acid (0.0005 mol), Pd(PPh<sub>3</sub>)<sub>4</sub> (0.00002 mol) and Na<sub>2</sub>CO<sub>3</sub> (0.0007 mol) in DME (3 ml) was stirred and refluxed in a sealed tube for 3 hours. H<sub>2</sub>O was added. The mixture was extracted with ethyl acetate. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 98/2; 15-40μm). The pure fractions were collected and the solvent was evaporated. Yield: 0.04 g of compound 21 (40%).

#### Example B12

The preparation of compound 23




A mixture of compound 22 (see table 3) (prepared according to B4.a) (0.0002 mol) and Raney Nickel (0.1 g) in CH<sub>3</sub>OH (10 ml) was stirred at room temperature for 15 minutes under a 2 bar pressure of H<sub>2</sub>, then filtered over celite. Celite was washed with CH<sub>3</sub>OH. The filtrate was evaporated. Yield: 0.48 g. This fraction was purified by column chromatography over kromasyl (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 15-40μm). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.13 g F1 and 0.13 g F2. F2 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.09 g of compound 23 (20%).



-87-

Example B13


The preparation of compound 24



A mixture of compound 1 (0.0004 mol) and Pd/C (0.07 g) in CH<sub>3</sub>OH (10 ml) was hydrogenated at room temperature for 5 hours under a 3 bar pressure of H<sub>2</sub>, then filtered over celite, washed with CH<sub>2</sub>Cl<sub>2</sub> and the solvent was evaporated till dryness. The residue (0.7 g) was purified by column chromatography over kromasyl (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0 to 99/1; 5 $\mu$ m). The pure fractions were collected and the solvent was evaporated. The residue (0.06 g) was crystallized from DIPE. The precipitate was filtered off and dried. Yield: 0.04 g of compound 24 (27%).

Example B14

The preparation of compound 26




NaH 60% (0.0004 mol) was added at room temperature to a mixture of compound 25 (see Table 4) (prepared according to B6.c) (0.0004 mol) in THF (30 ml). The mixture was stirred at room temperature for 1 hour. A solution of ICH<sub>3</sub> (0.0004 mol) in THF (30 ml) was added. The mixture was stirred at 60°C for 2 hours, then cooled, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (0.12 g) was purified by column chromatography over kromasil (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 10 $\mu$ m). The pure fractions were collected and the solvent was evaporated. Yield: 0.049 g of compound 26 (32%).

-88-


Example B15

a) The preparation of compound 123



Jones's reagent (0.0056 mol) was added at 5°C to a mixture of compound 18 (prepared according to B9) (0.0029 mol) in 2-propanone (20 ml) under N<sub>2</sub> flow. The mixture was stirred at 5°C for 2 hours, then poured out into H<sub>2</sub>O, basified with NaHCO<sub>3</sub> and  
5 extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (1.5 g) was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 98/2/0.1; 15-40μm). Two fractions (F1, F2) were collected and the solvent was evaporated. Yield: 0.122 g F1 (11%) and 0.19 g F2 (17%). F2 was crystallized from DIPE. The precipitate was  
10 filtered off and dried. Yield: 0.034 g of compound 123 (mp. 150°C).


b) The preparation of compound 124



A mixture of compound 123 (0.0005 mol) in POCl<sub>3</sub> (1.5 ml) was stirred at 80°C for 24 hours, poured out into ice and K<sub>2</sub>CO<sub>3</sub> 10% and extracted with CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered and the solvent was evaporated. The residue (0.14 g) was purified by column chromatography over kromasil (eluent:  
15 CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 99/1; 10μm). The pure fractions were collected and the solvent was evaporated. Yield: 0.026 g of compound 124.


Example B16

## a) The preparation of compound 125



NaOH 5N (2 ml) was added dropwise at 50°C to a mixture of compound 104 (see Table 3) (prepared according to B2.c) (0.0003 mol) and NH<sub>2</sub>OH, HCl (0.0004 mol) in ethanol (10 ml). The mixture was stirred at 50°C for 2 hours. Two-third of the mixture was evaporated. The mixture was poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with K<sub>2</sub>CO<sub>3</sub> 10%, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. Yield: 0.21 g of compound 125.

## b) The preparation of compound 126




1,1'-carbonyldiimidazole (0.0012 mol) was added to a mixture of compound 125 (0.0003 mol) in THF (20 ml). The mixture was stirred and refluxed overnight, poured out into H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (MgSO<sub>4</sub>), filtered, and the solvent was evaporated. The residue (0.17 g) was purified by column chromatography over kromasil (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 98/2; 10μm). Two fractions were collected and the solvent was evaporated. Yield: 0.035 g of fraction 1 and 0.05 g of fraction 2. Both fractions were mixed and crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.05 g of compound 126 (38%) (mp. >260°C).

-90-

Example B17

## Preparation of compound 253




(E)

- 5 a) 2.53 ml of acetonitrile, 0.056 g (0.253 mmol) of Pd(OAc)<sub>2</sub> and 0.154 g (0.506 mmol) of tris(2-methylphenyl)phosphine were brought in a 100 ml flask under nitrogen and the mixture was stirred for 10 minutes. To the mixture was added 1 g (2.53 mmol) of intermediate 58, 0.51 ml (3.8 mmol) of *N,N*-diethylethanamine and 0.36 g (5.06 mmol) of acrylamide. The mixture was heated at reflux (80°C) for 5 days yielding 28 % of compound 253.
- 10 b) In a 100ml flask under N<sub>2</sub> were introduced 0,8g (4,33mmol; 1 eq.) of intermediate 3a (E), 1g (4,33 mmol; 1 eq.) of intermediate 5 and 16ml of 2-propanol. To this mixture 0,72ml of HCl 6N in 2-propanol were added. The mixture was stirred under refluxed for 72 hours and then cooled yielding the hydrochloric acid salt of compound 253, i.e. compound 254.
- 15 Compound 254 can be converted into the free base according to art-known methodologies (see also Example B1A).

Compound 253 can be converted into compound 1 according to the method described above in Example A1c)y).
















The following Tables 3, 4 and 5 list compounds of formula (I) as prepared according to one of the above examples (Ex. No.).
















Table 3

















5










Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp. °C / (MH <sup>+</sup> )*
2	B2a	2-benzofuranyl	H	mp. > 240
21	B11	3-thienyl	H	mp. 220
3	B2b	2-furanyl	H	mp. 228
28	B2a	2-thienyl	H	mp. 235
29	B2a	phenyl	H	mp. 230
1	B1/B6a	-CH=CH-CN	H	mp. 245, (E)
30	B2a	2,4-dichlorophenyl	H	(460)
31	B2a	2-benzo[b]thienyl	H	(448)
32	B2a	1-naphthalenyl	H	(442)
33	B2a	3-chlorophenyl	H	(426)
34	B2a	3-acetylphenyl	H	(434)
35	B2a	3-methylphenyl	H	(406)
36	B2a	2-naphthalenyl	H	(442)
37	B2a	4-chlorophenyl	H	(426)
38	B2a	4-methoxyphenyl	H	(422)
39	B2a	4-methylthiophenyl	H	(438)
40	B2a		H	
19	B1		H	mp. 220
8	B5a	-C(=N-OH)-CH(CH <sub>3</sub> ) <sub>2</sub>	H	mp. 156
20	B10		H	mp. 205
27	B1		H	mp. 193







Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp. °C / (MH+)*
41	B10		H	mp. 200
42	B5a		H	mp. 155
43	B4b		H	mp. 110
44	B5b		H	mp. 110
45	B5a	-C(=N-OH)-CH <sub>3</sub>	H	mp. 135
9	B5b	-C(=N-O-CH <sub>3</sub> )-CH(CH <sub>3</sub> ) <sub>2</sub>	H	mp. 185
46	B5b		H	mp. 164
47	B4b	-CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	mp. 150
48	B4b		H	mp. 85
15	B6e		H	(461)
49	B6e		H	(449)
50	B6e		H	(487)
51	B6e		H	(493)
52	B6e		H	(473)
53	B6e		H	(443)
54	B6e		H	(446)
55	B6e		H	(449)
56	B6e		H	(521)











Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp. °C / (MH <sup>+</sup> )*
57	B6e		H	(457)
6	B4b	$-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_3)_2$	H	(430)
58	B4b		H	(506)
59	B4b		H	(428)
60	B4b		H	(532)
61	B4b		H	(504)
62	B4b		H	(503)
63	B4b		H	(472)
64	B4b		H	(491)
65	B4b	$-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$	H	(415)
66	B4b		H	(442)
67	B4b		H	(410)
68	B4b	$-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{CH}_3$	H	(401)
69	B4b		H	(399)
70	B4b		H	(396)
71	B4b	$-\text{CH}_2-\text{N}(\text{CH}_2-\text{CH}_2-\text{O}-\text{CH}_3)_2$	H	(461)
72	B4b		H	(485)
73	B4b		H	(456)
74	B4b		H	(492)
75	B4b	$-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{CN}$	H	(412)

Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp.°C / (MH+)*
76	B4b		H	(443)
77	B4b		H	(397)
78	B4b		H	(417)
79	B4b		H	(464)
80	B4b	-CH <sub>2</sub> -NH-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	mp, 105
81	B1		H	mp. 240
82	B10		H	mp. 170
24	B13	-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	mp. 208
83	B8		H	mp.>250°C
14	B6d		H	mp.158
84	B6c	-C(CH <sub>3</sub> )=CH-CN	H	mp.224°C (E)
18	B9	-CH(OH)-CH <sub>2</sub> -CN	H	mp.252°C
85	B4b		H	(474)
86	B4b		H	(473)
87	B4b		H	(426)
88	B4b		H	(424)
89	B4b		H	(446)
90	B4b		H	(397)



Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp.°C / (MH+)*
91	B4b		H	(438)
92	B4b		H	(438)
93	B4b		H	(410)
94	B4b		H	(410)
95	B4b		H	(478)
96	B4b		H	(473)
103	B6b	-CH=C(CH <sub>3</sub> )-CN	H	mp. 201°C (E)
11	B6b	-CH=C(CH <sub>3</sub> )-CN	H	mp. 246°C (Z)
10	B6a	-CH=CH-CN	H	mp. 258°C (Z)
4	B3	-CH <sub>2</sub> -CN	H	
17	B8		H	mp.110°C
97	B8		H	mp.240°C
16	B7		H	mp.>250°C
7	B4c	-CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	mp>260
5	B4a	4-thiomorpholinyl	-NO <sub>2</sub>	mp. 268
98	B4a	4-morpholinyl	-NO <sub>2</sub>	mp. 210
22	B4a	1-piperidiny1	-NO <sub>2</sub>	mp. 252
23	B12	1-piperidiny1	-NH <sub>2</sub>	mp. 262
12	B6c	H	-C(CH <sub>3</sub> )=CH-CN	(E) (381)
13	B6c	H	-C(CH <sub>3</sub> )=CH-CN	(Z) (381)
127	B1	-N(CH <sub>3</sub> ) <sub>2</sub>	H	mp. 228°C
123	B15a	-C(=O)-CH <sub>2</sub> -CN	H	mp. 150°C


Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp. °C / (MH+)*
116	B9C		H	(463)
128	B9C		H	(480)
129	B9C		H	(452)
130	B9C	-CH=N-NH-C(=O)-CH <sub>3</sub>	H	(400)
131	B9C	-CH=N-NH-C(=O)-CH <sub>2</sub> -CN	H	(425)
132	B9C		H	(468)
115	B9Bd	-C(=O)-NH-CH <sub>3</sub>	H	(373)
134	B9Bd	-C(=O)-N(CH <sub>3</sub> ) <sub>2</sub>	H	(387)
135	B9Bd	-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>3</sub>	H	(401)
136	B9Bd	-C(=O)-N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	(415)
137	B9Bd	-C(=O)-NH-CH <sub>2</sub> -CH <sub>3</sub>	H	(387)
138	B9Bd	-C(=O)-NH-CH <sub>2</sub> -CN	H	(398)
139	B9Bd	-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CN	H	(412)
140	B9Bd	-C(=O)-NH-CH <sub>2</sub> -C≡CH	H	(397)
141	B9Bd	-C(=O)-NH-CH <sub>2</sub> -CH=CH <sub>2</sub>	H	(399)
142	B9Bd	-C(=O)-NH-CH(CH <sub>3</sub> ) <sub>2</sub>	H	(401)
143	B1	-N[CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub> ] <sub>2</sub>	H	mp. 238°C
144	B13	-CH <sub>2</sub> -CH(CN) <sub>2</sub>	H	mp. 160°C
106	B6f	-CH=C(CN)-C(=O)-C(CH <sub>3</sub> ) <sub>3</sub>	H	(E), mp. 193°C
145	B9F		H	(E), mp. 229°C
146	B9F		H	(Z), mp. 258°C
147	B9Ea	-CH=C(CN)-CH <sub>2</sub> -CN	H	(Z/E=88/12) (406)
148	B6c	-C(CH <sub>2</sub> -CH <sub>3</sub> )=CH-CN	H	(E), mp. 173°C
149	B6c	-C(CH(CH <sub>3</sub> ) <sub>2</sub> )=CH-CN	H	(E), mp. 132°C
150	B6c	-C(CH(CH <sub>3</sub> ) <sub>2</sub> )=CH-CN	H	(Z), mp. 132°C
151	B6b	-CH=C(CH <sub>3</sub> )-CN	H	(Z), mp. 246°C

Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp.°C / (MH+)*
152	B6b	-CH=C(CH <sub>3</sub> )-CN	H	(E), mp. 201°C
153	B13	-CH <sub>2</sub> -CH(CH <sub>3</sub> )-CN	H	mp. 187°C
124	B15b	-C(Cl)=CH-CN	H	
154	B9Ba	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CN	H	(E)
112	B9Ba	-CH=CH-C(=O)-N(CH <sub>3</sub> ) <sub>2</sub>	H	(E), mp.>264°C
155	B9Bc		H	(E), mp. 156°C
156	B9Bc		H	(E), mp. 168°C
157	B9Bc		H	(E), mp.>265°C
158	B9Bc	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>3</sub>	H	(E), mp.>260°C
114	B9Bc	-CH=CH-C(=O)-N(CH <sub>3</sub> )-(CH <sub>2</sub> ) <sub>2</sub> -CN	H	(E), mp. 168°C
159	B9Bc	-CH=CH-C(=O)-N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	(E), mp. 249°C
160	B6b	-C(CH <sub>3</sub> )=C(CH <sub>3</sub> )-CN	H	(E)
107	B9Aa	-CH=CH-Cl	H	(Z), mp. 250°C
161	B9Aa	-CH=CH-Br	H	(Z), mp. 248°C
111	B9Ad	-CH=C(Br) <sub>2</sub>	H	mp. 223°C
122	B9F		H	(E), mp. 120°C
162	B9F		H	(E), mp. >260°C
163	B9F		H	mp. 128°C
164	B9FF		H	mp. 104°C
125	B16a		H	
104	B2c		H	
165	B9F		H	mp. 112°C









Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4</sup>	Physical data mp. °C / (MH <sup>+</sup> )*
166	B9F		H	mp. 194°C
167	B9F		H	mp. 191°C
126	B16b		H	mp. >260°C
168	B4c	-CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>3</sub>	H	mp. 201°C
117	B9Da	H	-N(CH <sub>3</sub> ) <sub>2</sub>	mp. 132°C
120	B9Eb	-CH=C(CN) <sub>2</sub>	H	
253	B17a/b	-CH=CH-C(=O)NH <sub>2</sub>	H	(E)
254	B17b	-CH=CH-C(=O)NH <sub>2</sub>	H	(E) HCl








\* (MH<sup>+</sup>) defines the mass of the protonated compound; it was determined with a MicroMass spectrometer equipped with an electrospray probe with a quadripolar analyser.



5 Table 4:



Comp No.	Ex. No.	R <sup>3</sup>	R <sup>1</sup>	Physical data mp. °C / (MH <sup>+</sup> )*
25	B6c	-CH=CH-CN	H	mp. 256°C
99	B3	-CH <sub>2</sub> -CN	H	mp. 184°C
100	B4b	-CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	mp. 172 °C
102	B13	-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	mp. 224°C
101	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	mp. 196°C
26	B14	-CH=CH-CN	CH <sub>3</sub>	mp. 195°C
169	B9Bd	-C(=O)-N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	mp. 172°C
170	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CN	H	
171	B4b		H	(398)
172	B2a		H	mp. 158°C


Comp No.	Ex. No.	R <sup>3</sup>	R <sup>1</sup>	Physical data mp. °C / (MH+)*
173	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>3</sub> ) <sub>2</sub>	H	mp. 196°C
174	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH=N-CN	H	mp. 254°C
175	B14	2-furanyl	CH <sub>3</sub>	mp. 178°C
118	B9Db		H	164°C
176	B14		CH <sub>3</sub>	mp. 188°C
177	B9Aa	-CH=CH-Br	H	(Z), mp. 169°C
110	B9Ac	-CH=C(F)-CN	H	(E), mp. 254°C
178	B6b	-CH=C(CH <sub>3</sub> )-CN	H	(Z)
179	B6b	-CH=C(CH <sub>3</sub> )-CN	H	(E)
180	B9Bb		H	(E)
181	B9Bc	-CH=CH-C(=O)-NH-cyclopropyl	H	(E) (426)
182	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>3</sub> ) <sub>2</sub>	H	(E) (427)
183	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -O-CH <sub>3</sub>	H	(E)(458)
184	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	H	(E)(442)
185	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	(E)439)
186	B9Bc		H	(E)(468)
187	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>3</sub> ) <sub>2</sub>	H	(E)(471)
188	B9Bc	-CH=CH-C(=O)-NH-(CH <sub>2</sub> ) <sub>3</sub> -O-CH <sub>2</sub> -CH <sub>3</sub>	H	(E)(472)
189	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH <sub>3</sub>	H	(E)(414)
190	B9Bc	-CH=CH-C(=O)-NH-CH <sub>2</sub> -CH <sub>2</sub> -O-CH <sub>3</sub>	H	(E)(444)
191	B9Bc	-CH=CH-C(=O)-NH-CH(CH <sub>3</sub> ) <sub>2</sub>	H	(E)(428)
192	B4b		H	(E)(491)
193	B4b		H	(E)(444)
194	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	(E)(439)
195			H	(E)(483)
196	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>2</sub> -O-CH <sub>3</sub> ) <sub>2</sub>	H	(E)(488)
197	B4b		H	(E)(476)
198	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>3</sub>	H	(E)(428)

Comp No.	Ex. No.	R <sup>3</sup>	R <sup>1</sup>	Physical data mp.°C / (MH+)*
199	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	(E)(485)
200	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>3</sub> )-CH <sub>3</sub>	H	(E)(414)
201	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	(E)(456)
202	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>3</sub>	H	(E)(442)
203	B4b		H	(E)(438)
204	B4b		H	(E)(442)
205	B4b		H	(E)(455)
206	B4b	-CH=CH-CH <sub>2</sub> -N(benzyl)-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>3</sub> ) <sub>2</sub>	H	(E)(533)
207	B4b	-CH=CH-CH <sub>2</sub> -N(CH <sub>3</sub> ) <sub>2</sub>	H	(E)(457)
208	B4b	-CH=CH-CH <sub>2</sub> -N(isopropyl) <sub>2</sub>	H	(E)(456)
121	B9Bb	-CH=CH-C(=O)-NH <sub>2</sub>	H	(E)
209	B9Bb		H	(E), mp. 116°C
210	B9Bb		H	(E), mp. 254°C
211	B9Bb	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -OH	H	(E), mp. 222°C
212	B9Ba	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CN	H	(E), mp. 198°C
213	B6c	-C(CH <sub>3</sub> )=CH-CN	H	(E)
214	B9Bc	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	(E), mp. 204°C
215	B9Bc	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>3</sub>	H	(E), mp. 211°C
216	B9Bc		H	(E), mp. 246°C
217	B9Bc	-CH=CH-C(=O)-N(CH <sub>2</sub> -CH <sub>3</sub> ) <sub>2</sub>	H	(E), mp. 226°C
218	B9Bc		H	(E), mp. 196°C
219	B9Ba	-CH=CH-C(=O)-N(CH <sub>3</sub> ) <sub>2</sub>	H	(E), mp. 225°C
220	B9E	-CH=C(CN)-CH <sub>2</sub> -CN	H	(Z), mp. 195°C
109	B9Ab	-CH=CH-Cl	H	(E), mp. 200°C
108	B9Ab	-CH=CH-Cl	H	(Z), mp. 165°C
221	B9Ba	-CH=CH-C(=O)-NH-CH <sub>3</sub>	H	(E), mp. 260°C
222	B9Bb	-CH=CH-C(=O)-N(CH <sub>2</sub> -CH <sub>2</sub> -O-CH <sub>3</sub> ) <sub>2</sub>	H	(E), mp. 158°C


Comp No.	Ex. No.	R <sup>3</sup>	R <sup>1</sup>	Physical data mp.°C / (MH <sup>+</sup> )*
223	B9Bb		H	(E), mp. 208°C
224	B9Bb		H	(E), mp. 208°C
113	B9Bb	-CH=CH-C(=O)-N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>3</sub>	H	(E), mp. 212°C
225	B4b	-CH <sub>2</sub> -N(CH <sub>2</sub> -CH <sub>2</sub> -CN) <sub>2</sub>	H	mp. 154°C
226	B2a	2-furanyl	H	mp. 162°C

\* (MH<sup>+</sup>) defines the mass of the protonated compound; it was determined with a MicroMass spectrometer equipped with an electrospray probe with a quadripolar analyser.

5 Table 5:



Comp No.	Ex. No.	R <sup>3</sup>	R <sup>4a</sup>	R <sup>4b</sup>	X <sup>1</sup>	Physical data mp.°C
227	B13	-CH <sub>2</sub> -CH <sub>2</sub> -CN	CH <sub>3</sub>	H	-NH	mp. 186°C
228	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CN	CH <sub>3</sub>	H	-NH	mp. 138°C
229	B6b	-CH=C(CH <sub>3</sub> )-CN	CH <sub>3</sub>	H	-NH	mp. 190°C
230	B6c	-CH=CH-CN	CH <sub>3</sub>	H	-O-	(E), mp. 254°C
231	B6b	-CH=C(CH <sub>3</sub> )-CN	CH <sub>3</sub>	H	-O-	mp. 150°C
232	B6c	-C(CH <sub>3</sub> )=CH-CN	CH <sub>3</sub>	H	-O-	(E), mp. 234°C
105	B4d	-CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>3</sub>	CH <sub>3</sub>	H	-O-	mp. 140°C
233	B6b	-CH=C(CH <sub>3</sub> )-CN	CH <sub>3</sub>	Cl	-NH	mp. 214°C
234	B13	-CH <sub>2</sub> -CH <sub>2</sub> -CN	CH <sub>3</sub>	H	-O-	mp. 199°C
235	B13	-CH(CH <sub>3</sub> )-CH <sub>2</sub> -CN	CH <sub>3</sub>	H	-O-	mp. 195°C
236	B13	-CH <sub>2</sub> -CH(CH <sub>3</sub> )-CN	CH <sub>3</sub>	H	-O-	mp. 161°C
237	B6c	-CH=CH-CN	CH <sub>3</sub>	H	-NH	(E), mp. >264°C
238	B3	-CH <sub>2</sub> -CN	CH <sub>3</sub>	Cl	-NH	mp. 184°C
239	B6c	-CH=CH-CN	CH <sub>3</sub>	2-furanyl	-NH	(E) mp. 175°C
119	B9E	-CH=C(CN)-CH <sub>2</sub> -CN	CH <sub>3</sub>	2-furanyl	-NH	

Comp p No.	Ex. No.	R <sup>3</sup>	R <sup>4a</sup>	R <sup>4b</sup>	X <sup>1</sup>	Physical data mp. °C
240	B9F		CH <sub>3</sub>	Cl	-NH	mp. 248°C Z/E=50/50
241	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CN	CH <sub>3</sub>	Br	-NH	mp. 148°C
242	B1	-CH=CH-CN	H	isopropyl	-NH	(E) 30%-(Z) 70%
243	B4b	-CH <sub>2</sub> -N(CH <sub>3</sub> )-CH <sub>2</sub> -CH <sub>2</sub> -CN	CH <sub>3</sub>	Cl	-NH	mp. 85°C
244	B6c	-CH=CH-CN	H	Br	-NH	(E), mp. 270°C
245	B6c	-CH=CH-CN	H	-OCH <sub>3</sub>	-NH	(E), mp. 258°C
246	B6b	-C(CH <sub>3</sub> )=C(CH <sub>3</sub> )-CN	CH <sub>3</sub>	H	-O-	(E), mp. 214°C
247	B6b	-CH=C(CH <sub>3</sub> )-CN	CH <sub>3</sub>	Br	-NH	mp. 212°C
248	B6c	-CH=CH-CN	CH <sub>3</sub>	Br	-NH	(E), mp. 250°C
249	B6b	-CH=C(CH <sub>3</sub> )-CN	H	-OCH <sub>3</sub>	-NH	mp. 166°C
250	B6b	-CH=C(CH <sub>3</sub> )-CN	H	Br	-NH	mp. 186°C
251	B13	-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	-OCH <sub>3</sub>	-NH	mp. 228°C
252	B4c	-CH <sub>2</sub> -O-CH <sub>2</sub> -CH <sub>2</sub> -CN	H	Cl	-NH	mp. 168°C
133	B6c	-CH=CH-CN	CH <sub>3</sub>	Cl	-NH	(E), mp, 258°C

### C. Pharmacological example

The pharmacological activity of the present compounds was examined using the following test.

- 5 A rapid, sensitive and automated assay procedure was used for the *in vitro* evaluation of anti-HIV agents. An HIV-1 transformed T4-cell line, MT-4, which was previously shown (Koyanagi et al., *Int. J. Cancer*, **36**, 445-451, 1985) to be highly susceptible to and permissive for HIV infection, served as the target cell line. Inhibition of the HIV-induced cytopathic effect was used as the end point. The viability of both HIV- and
- 10 mock-infected cells was assessed spectrophotometrically via the *in situ* reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT). The 50% cytotoxic concentration (CC<sub>50</sub> in M) was defined as the concentration of compound that reduced the absorbance of the mock-infected control sample by 50%. The percent protection achieved by the compound in HIV-infected cells was calculated by the
- 15 following formula :

$$\frac{(\text{OD}_T)_{\text{HIV}} - (\text{OD}_C)_{\text{HIV}}}{(\text{OD}_C)_{\text{MOCK}} - (\text{OD}_C)_{\text{HIV}}} \quad \text{expressed in \%},$$



-103-

whereby  $(OD_T)_{HIV}$  is the optical density measured with a given concentration of the test compound in HIV-infected cells;  $(OD_C)_{HIV}$  is the optical density measured for the control untreated HIV-infected cells;  $(OD_C)_{MOCK}$  is the optical density measured for the control untreated mock-infected cells; all optical density values were determined at 540 nm. The dose achieving 50% protection according to the above formula was defined as the 50% inhibitory concentration ( $IC_{50}$  in M). The ratio of  $CC_{50}$  to  $IC_{50}$  was defined as the selectivity index (SI).

Table 6 lists the  $pIC_{50}$  ( $-\log IC_{50}$ ),  $pCC_{50}$  ( $-\log CC_{50}$ ) and  $pSI$  ( $pCC_{50} - pIC_{50}$ ) values for the compounds of formula (I). For example, a compound with a  $IC_{50}$  value of  $10^{-9}M$ , i.e.  $pIC_{50}=9$ , and a  $CC_{50}$  value of  $10^{-5}M$ , i.e.  $pCC_{50}=5$ , has a SI of  $10^{-5}M/10^{-9}M = 10.000$ , i.e. a  $pSI$  of  $5-9 = -4$ .

Table 6

Co. No.	$pIC_{50}$ (M)	$pCC_{50}$ (M)	$pSI$
21	8.4	4.9	-3.5
3	8.4	5.5	-2.9
1	9.4	5.0	-4.4
34	8.0	4.8	-3.2
19	8.4	4.8	-3.6
45	8.7	5.0	-3.8
49	8.0	4.8	-3.2
70	8.1	4.8	-3.3
75	9.0	5.0	-4.0
78	8.4	4.9	-3.5
79	8.0	5.3	-2.7
84	9.0	4.5	-4.5
18	8.8	4.9	-4.0
25	9	4	-5
24	9.1	5.7	-3.4
81	9.1	5.6	-3.5
11	9.2	5.7	-3.5
10	9.2	6.3	-2.9
174	8.8	5.3	-3.5
227	9.5	<4.0	<-5.5
144	8.6	6.4	-2.2

-104-

Co. No.	pIC <sub>50</sub> (M)	pCC <sub>50</sub> (M)	pSI
229	8.8	<4.0	<-4.8
118	8.4	4.1	<-4.1
177	8.3	<4.0	<-4.3
106	7.7	5.2	-2.5
145	8.7	5.3	-3.4
147	9.4	5.7	-3.7
148	8.8	4.9	-3.9
230	9.2	<4.0	<-5.2
231	9.2	<4.0	<-5.2
232	8.4	<4.0	<-4.4
105	7.2	<4.0	<-3.2
110	8.6	4.3	-4.3
233	9.3	5.7	-3.6
234	8.7	<4.0	<-4.7
235	9.3	<4.0	<-5.3
236	8.8	<4.0	<-4.8
149	9.1	5.3	-3.8
150	8.8	4.8	-4.0
237	8.9	<4.0	<-4.9
151	9.1	5.5	-3.6
152	9.1	4.8	-4.3
178	8.8	5.7	-3.1
179	8.9	<4.0	<-4.9
153	9.2	6.3	-2.9
124	8.5	4.7	-3.8
238	9.5	5.6	-3.9
112	9.1	4.9	-4.2
244	9.2	4	-5.2
209	8.6	4.9	-3.7
210	8.3	4.8	-3.5
155	8.8	6.3	-2.5
156	7.7	5.1	-2.6
158	8	5.5	-2.5
212	9.1	5	-4.1
114	8.6	5.1	-3.5


-105-

Co. No.	pIC50 (M)	pCC50 (M)	pSI
213	9	4.8	-4.2
214	8.6	5.1	-3.5
215	9.1	5.5	-3.6
216	8.2	5	-3.6
219	9.1	5	-4.1
245	8.8	4	-4.8
146	8.4	5.4	-3
247	9.2	6.2	-3
248	9.3	5.7	-3.5
249	8.5	4	-4.5
42	9	6.3	-2.7
251	8.9	5	-3.9
133	9.2	4	-5.2
9	8.8	4.8	-4
239	8.9	5	-3.9
241	9.4	5.3	-4.1
126	8.4	4.9	-3.5

-106-

Claims


1. A compound of formula

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a

5 stereochemically isomeric form thereof, wherein

 $-a^1=a^2-a^3=a^4-$  represents a bivalent radical of formula $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$  (a-1); $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$  (a-2); $-\text{N}=\text{CH}-\text{N}=\text{CH}-$  (a-3);10  $-\text{N}=\text{CH}-\text{CH}=\text{N}-$  (a-4); $-\text{N}=\text{N}-\text{CH}=\text{CH}-$  (a-5); $-b^1=b^2-b^3=b^4-$  represents a bivalent radical of formula $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$  (b-1); $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$  (b-2);15  $-\text{N}=\text{CH}-\text{N}=\text{CH}-$  (b-3); $-\text{N}=\text{CH}-\text{CH}=\text{N}-$  (b-4); $-\text{N}=\text{N}-\text{CH}=\text{CH}-$  (b-5); $n$  is 0, 1, 2, 3 or 4; and in case  $-a^1=a^2-a^3=a^4-$  is (a-1), then  $n$  may also be 5; $m$  is 1, 2, 3 and in case  $-b^1=b^2-b^3=b^4-$  is (b-1), then  $m$  may also be 4;20  $R^1$  is hydrogen; aryl; formyl;  $C_{1-6}$ alkylcarbonyl;  $C_{1-6}$ alkyl;  $C_{1-6}$ alkyloxycarbonyl; $C_{1-6}$ alkyl substituted with formyl,  $C_{1-6}$ alkylcarbonyl,  $C_{1-6}$ alkyloxycarbonyl, $C_{1-6}$ alkylcarbonyloxy;  $C_{1-6}$ alkyloxy $C_{1-6}$ alkylcarbonyl substituted with $C_{1-6}$ alkyloxycarbonyl;each  $R^2$  independently is hydroxy, halo,  $C_{1-6}$ alkyl optionally substituted with cyano or25  $-\text{C}(=\text{O})\text{R}^6$ ,  $C_{3-7}$ cycloalkyl,  $C_{2-6}$ alkenyl optionally substituted with one or more halogen atoms or cyano,  $C_{2-6}$ alkynyl optionally substituted with one or more halogen atoms or cyano,  $C_{1-6}$ alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di( $C_{1-6}$ alkyl)amino, polyhalomethyl, polyhalomethylthio,  $-\text{S}(=\text{O})_p\text{R}^6$ ,  $-\text{NH}-\text{S}(=\text{O})_p\text{R}^6$ ,  $-\text{C}(=\text{O})\text{R}^6$ ,  $-\text{NHC}(=\text{O})\text{H}$ ,  $-\text{C}(=\text{O})\text{NHNH}_2$ ,  $-\text{NHC}(=\text{O})\text{R}^6$ ,  $-\text{C}(=\text{NH})\text{R}^6$  or a radical of

30 formula

wherein each  $A_1$  independently is N, CH or  $\text{CR}^6$ ; and

-107-

A<sub>2</sub> is NH, O, S or NR<sup>6</sup>;

X<sub>1</sub> is -NR<sup>5</sup>-, -NH-NH-, -N=N-, -O-, -C(=O)-, C<sub>1-4</sub>alkanediyl, -CHOH-, -S-, -S(=O)<sub>p</sub>-,  
-X<sub>2</sub>-C<sub>1-4</sub>alkanediyl- or -C<sub>1-4</sub>alkanediyl-X<sub>2</sub>-;

X<sub>2</sub> is -NR<sup>5</sup>-, -NH-NH-, -N=N-, -O-, -C(=O)-, -CHOH-, -S-, -S(=O)<sub>p</sub>-;

- 5 R<sup>3</sup> is NHR<sup>13</sup>; NR<sup>13</sup>R<sup>14</sup>; -C(=O)-NHR<sup>13</sup>; -C(=O)-NR<sup>13</sup>R<sup>14</sup>; -C(=O)-R<sup>15</sup>; -CH=N-NH-  
C(=O)-R<sup>16</sup>; C<sub>1-6</sub>alkyl substituted with one or more substituents each  
independently selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or  
R<sup>7</sup>; C<sub>1-6</sub>alkyl substituted with one or more substituents each independently  
selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup> and  
10 wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  
C<sub>1-4</sub>alkanediyl; C<sub>1-6</sub>alkyl substituted with hydroxy and a second substituent  
selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>;  
C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl optionally substituted with one or more substituents each  
independently selected from cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or  
15 R<sup>7</sup>; C<sub>2-6</sub>alkenyl substituted with one or more substituents each independently  
selected from halo, cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>;  
C<sub>2-6</sub>alkynyl substituted with one or more substituents each independently  
selected from halo, cyano, NR<sup>9</sup>R<sup>10</sup>, -C(=O)-NR<sup>9</sup>R<sup>10</sup>, -C(=O)-C<sub>1-6</sub>alkyl or R<sup>7</sup>;  
-C(=N-O-R<sup>8</sup>)-C<sub>1-4</sub>alkyl; R<sup>7</sup> or -X<sub>3</sub>-R<sup>7</sup>;
- 20 X<sub>3</sub> is -NR<sup>5</sup>-, -NH-NH-, -N=N-, -O-, -C(=O)-, -S-, -S(=O)<sub>p</sub>-, -X<sub>2</sub>-C<sub>1-4</sub>alkanediyl-,  
-C<sub>1-4</sub>alkanediyl-X<sub>2a</sub>-, -C<sub>1-4</sub>alkanediyl-X<sub>2b</sub>-C<sub>1-4</sub>alkanediyl,  
-C(=N-OR<sup>8</sup>)-C<sub>1-4</sub>alkanediyl-;

with X<sub>2a</sub> being -NH-NH-, -N=N-, -O-, -C(=O)-, -S-, -S(=O)<sub>p</sub>-; and

with X<sub>2b</sub> being -NH-NH-, -N=N-, -C(=O)-, -S-, -S(=O)<sub>p</sub>-;

- 25 R<sup>4</sup> is halo, hydroxy, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, cyano, nitro,  
polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl, C<sub>1-6</sub>alkyloxycarbonyl,  
C<sub>1-6</sub>alkylcarbonyl, formyl, amino, mono- or di(C<sub>1-4</sub>alkyl)amino or R<sup>7</sup>;

R<sup>5</sup> is hydrogen; aryl; formyl; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxycarbonyl;  
C<sub>1-6</sub>alkyl substituted with formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>1-6</sub>alkyloxycarbonyl or  
30 C<sub>1-6</sub>alkylcarbonyloxy; C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkylcarbonyl substituted with  
C<sub>1-6</sub>alkyloxycarbonyl;

R<sup>6</sup> is C<sub>1-4</sub>alkyl, amino, mono- or di(C<sub>1-4</sub>alkyl)amino or polyhaloC<sub>1-4</sub>alkyl;

- R<sup>7</sup> is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic  
carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or  
35 aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems  
may optionally be substituted with one, two, three, four or five substituents each  
independently selected from halo, hydroxy, mercapto, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl,

-108-

aminoC<sub>1-6</sub>alkyl, mono or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylthio, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl, -CH(=N-O-R<sup>8</sup>), R<sup>7a</sup>, -X<sub>3</sub>-R<sup>7a</sup> or R<sup>7a</sup>-C<sub>1-4</sub>alkyl;

- 5 R<sup>7a</sup> is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, aminoC<sub>1-6</sub>alkyl, mono or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, formyl, C<sub>1-6</sub>alkylcarbonyl, C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylthio, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl, -CH(=N-O-R<sup>8</sup>);

R<sup>8</sup> is hydrogen, C<sub>1-4</sub>alkyl, aryl or arylC<sub>1-4</sub>alkyl;

- R<sup>9</sup> and R<sup>10</sup> each independently are hydrogen; hydroxy; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl; -CH(=NR<sup>11</sup>) or R<sup>7</sup>, wherein each of the aforementioned C<sub>1-6</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, mono- or di(C<sub>1-4</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup>, R<sup>7</sup>; or

R<sup>9</sup> and R<sup>10</sup> may be taken together to form a bivalent or trivalent radical of formula

- 25 -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>- (d-1)  
 -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>- (d-2)  
 -CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>- (d-3)  
 -CH<sub>2</sub>-CH<sub>2</sub>-S-CH<sub>2</sub>-CH<sub>2</sub>- (d-4)  
 -CH<sub>2</sub>-CH<sub>2</sub>-NR<sup>12</sup>-CH<sub>2</sub>-CH<sub>2</sub>- (d-5)  
 -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>- (d-6)  
 30 =CH-CH=CH-CH=CH- (d-7)

R<sup>11</sup> is cyano; C<sub>1-4</sub>alkyl optionally substituted with C<sub>1-4</sub>alkyloxy, cyano, amino, mono- or di(C<sub>1-4</sub>alkyl)amino or aminocarbonyl; C<sub>1-4</sub>alkylcarbonyl; C<sub>1-4</sub>alkyloxycarbonyl; aminocarbonyl; mono- or di(C<sub>1-4</sub>alkyl)aminocarbonyl;

R<sup>12</sup> is hydrogen or C<sub>1-4</sub>alkyl;

- 35 R<sup>13</sup> and R<sup>14</sup> each independently are C<sub>1-6</sub>alkyl optionally substituted with cyano or aminocarbonyl, C<sub>2-6</sub>alkenyl optionally substituted with cyano or aminocarbonyl, C<sub>2-6</sub>alkynyl optionally substituted with cyano or aminocarbonyl;

-109-

$R^{15}$  is  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl;

$R^{16}$  is  $C_{1-6}$ alkyl optionally substituted with cyano or aminocarbonyl, or  $R^7$ ;


p is 1 or 2;

- aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl, mono or di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylthio, cyano, nitro, polyhalo $C_{1-6}$ alkyl, polyhalo $C_{1-6}$ alkyloxy, aminocarbonyl,  $R^7$  or  $-X_3-R^7$ .
2. A compound according to claim 1 wherein  $R^3$  is  $C_{1-6}$ alkyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{1-6}$ alkyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$  and wherein 2 hydrogen atoms bound at the same carbon atom are replaced by  $C_{1-4}$ alkanediyl;  $C_{1-6}$ alkyl substituted with hydroxy and a second substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{2-6}$ alkenyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $C_{2-6}$ alkynyl substituted with at least one substituent selected from cyano, aminocarbonyl,  $NR^9R^{10}$  or  $R^7$ ;  $-C(=N-O-R^8)-C_{1-4}$ alkyl;  $R^7$  or  $-X_3-R^7$ ;  $R^4$  is halo, hydroxy,  $C_{1-6}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy, cyano, nitro, polyhalo $C_{1-6}$ alkyl, polyhalo $C_{1-6}$ alkyloxy, aminocarbonyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylcarbonyl, formyl, amino, mono- or di( $C_{1-4}$ alkyl)amino;  $R^7$  is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl, mono or di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,  $C_{3-7}$ cycloalkyl,  $C_{1-6}$ alkyloxy,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkylthio, cyano, nitro, polyhalo $C_{1-6}$ alkyl, polyhalo $C_{1-6}$ alkyloxy, aminocarbonyl,  $R^{7a}$ ,  $-X_3-R^{7a}$  or  $R^{7a}-C_{1-4}$ alkyl;  $R^{7a}$  is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl, amino $C_{1-6}$ alkyl, mono or di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,

-110-

C<sub>3-7</sub>cycloalkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxycarbonyl, C<sub>1-6</sub>alkylthio, cyano, nitro, polyhaloC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyloxy, aminocarbonyl; R<sup>9</sup> and R<sup>10</sup> each independently are hydrogen; hydroxy; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyloxy; C<sub>1-6</sub>alkylcarbonyl; C<sub>1-6</sub>alkyloxycarbonyl; amino; mono- or di(C<sub>1-6</sub>alkyl)amino; mono- or di(C<sub>1-6</sub>alkyl)aminocarbonyl or R<sup>7</sup>, wherein each of the aforementioned C<sub>1-6</sub>alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyloxy, carboxyl, C<sub>1-6</sub>alkyloxycarbonyl, cyano, amino, imino, mono- or di(C<sub>1-4</sub>alkyl)amino, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)<sub>p</sub>R<sup>6</sup>, -NH-S(=O)<sub>p</sub>R<sup>6</sup>, -C(=O)R<sup>6</sup>, -NHC(=O)H, -C(=O)NHNH<sub>2</sub>, -NHC(=O)R<sup>6</sup>, -C(=NH)R<sup>6</sup>, R<sup>7</sup>.

3. A compound according to claim 1 or 2 having the formula



wherein


-a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>-, -b<sup>1</sup>=b<sup>2</sup>-b<sup>3</sup>=b<sup>4</sup>-, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, m and X<sub>1</sub> are as defined in claim 1;

n' is 0, 1, 2 or 3 and in case -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- is (a-1), then n' may also be 4;

R<sup>2'</sup> is halo, C<sub>1-6</sub>alkyl, trihalomethyl, trihalomethyloxy, cyano, aminocarbonyl, C<sub>1-6</sub>alkyl substituted with cyano or aminocarbonyl;

provided that R<sup>2'</sup> is placed at the para position in respect of the NR<sup>1</sup> moiety.

4. A compound according to any one of claims 1 to 3 having the formula



wherein


-b<sup>1</sup>=b<sup>2</sup>-b<sup>3</sup>=b<sup>4</sup>-, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, m and X<sub>1</sub> are as defined in claim 1;

n' and R<sup>2'</sup> are as defined in claim 3.

5. A compound according to any one of claims 1 to 4 having the formula



-111-



wherein

$R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $X_1$  are as defined in claim 1;

$n'$  and  $R^{2'}$  are as defined in claim 3.

5

6. A compound according to any one of claims 3 to 5 wherein  $R^{2'}$  is cyano, aminocarbonyl or  $C_{1-6}$ alkyl substituted with cyano or aminocarbonyl.

7. A compound according to any one of claims 1 to 6 for use as a medicine.

10

8. The use of a compound as defined in any one of claims 1 to 6 for the manufacture of a medicament for the prevention or the treatment of HIV (Human Immunodeficiency Virus) infection.

15

9. The use of a compound according to claim 8 for the manufacture of a medicament for the prevention or the treatment of drug resistant HIV infection.

10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 6.


20

11. A process for preparing a pharmaceutical composition according to claim 10 characterized in that a therapeutically effective amount of a compound as claimed in any one of claims 1 to 6 is intimately mixed with a pharmaceutically acceptable carrier.

25


12. A process for preparing a compound as claimed in claim 1, characterized by  
a) reacting an intermediate of formula (II) with an intermediate of formula (III)

-112-




with  $W_1$  being a suitable leaving group, and  $R^1, R^2, R^3, R^4, X_1, m, n, -a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;

- b) reacting an intermediate of formula (IV) with an intermediate of formula (V) in the presence of a suitable catalyst, a suitable salt and a suitable solvent



with  $W_2$  being a suitable leaving group, and  $R^1, R^2, R^4, X_1, m, n, -a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1 and  $R^{7'}$  representing a monocyclic, bicyclic or tricyclic aromatic ring system and  $R^a$  representing a boronate or a tri( $C_{1-4}$ alkyl)stannane;


- c) reacting an intermediate of formula (IV) with an intermediate of formula (VI)



with  $W_2$  being a suitable leaving group, and  $R^1, R^2, R^4, X_1, m, n, -a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1 and  $R^{7''}$  representing a monocyclic, bicyclic or tricyclic saturated ring system;

- d) reacting an intermediate of formula (VII) with a suitable cyanide salt in the presence of a suitable solvent


-113-



with  $W_3$  being a suitable leaving group, and  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;


e) reacting an intermediate of formula (VII) with an intermediate of formula (VIII)

5 optionally in the presence of a suitable salt and a suitable solvent



with  $W_3$  being a suitable leaving group, and  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1 and  $Q$  representing  $R^7$ ;  $NR^9R^{10}$  or  $C_{1-6}$ alkyloxy optionally substituted with  $CN$ ,  $R^7$  or  $NR^9R^{10}$ ;

10 f) reacting an intermediate of formula (IX) with an intermediate of formula (X) in the presence of a suitable solvent




with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $R^8$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;

g) reacting an intermediate of formula (XI) with a Wittig or Horner-Emmons reagents of formula (XII) representing a suitable precursor of a phosphorus ylide in the presence of a suitable salt and a suitable solvent


15

-114-




with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1,  $R^c$  representing hydrogen or  $C_{1-4}$ alkyl,  $R^{c'}$  representing hydrogen,  $C_{1-4}$ alkyl or  $R^7$ , provided that  $CR^{c'}=CR^c$  is limited to  $C_{2-6}$ alkenyl, and  $R^b$  representing (Phenyl) $_3P^+-Cl^-$  or  $(CH_3CH_2-O)_2P(=O)-$ ;

h) reacting an intermediate of formula (XI) with an intermediate of formula (XIII) in the presence of a suitable solvent



with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $R^7$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1,  $R^c$  representing hydrogen or  $C_{1-4}$ alkyl,  $R^{c'}$  representing  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;


i) reacting an intermediate of formula (XI-b) with 2-butenedinitrile in the presence of tributylphosphine and a suitable solvent




with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;

j) reacting an intermediate of formula (XI-b) with propanedinitrile in the presence of a suitable base and a suitable solvent

-115-




k) reacting an intermediate of formula (XI-b) with CH<sub>3</sub>-CN in the presence of a suitable proton abstracting agent, a suitable substrate for the proton abstracting agent and in the presence of a suitable solvent



5

with R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, X<sub>1</sub>, m, n, -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- and -b<sup>1</sup>=b<sup>2</sup>-b<sup>3</sup>=b<sup>4</sup>- as defined in claim 1;

l) reacting an intermediate of formula (XI) with a Wittig or Horner-Emmons reagents of formula (XII') representing a suitable precursor of a phosphorus ylide in the presence of nBuLi and a suitable solvent




10

with R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, X<sub>1</sub>, m, n, -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- and -b<sup>1</sup>=b<sup>2</sup>-b<sup>3</sup>=b<sup>4</sup>- as defined in claim 1, R<sup>c</sup> representing hydrogen or C<sub>1-4</sub>alkyl, R<sup>c'</sup> representing hydrogen, C<sub>1-4</sub>alkyl or R<sup>7</sup>, provided that CR<sup>c'</sup>=CR<sup>c</sup> is limited to C<sub>2-6</sub>alkenyl, and R<sup>b</sup> representing (Phenyl)<sub>3</sub>P<sup>+</sup>- Cl<sup>-</sup> or (CH<sub>3</sub>CH<sub>2</sub>-O)<sub>2</sub>P(=O)-;

15


m) reacting an intermediate of formula (XI-a) with an intermediate of formula (XIII') in the presence of a suitable Horner-Emmons reagent, nBuLi, 1,1,1-trimethyl-N-(trimethylsilyl)-silanamine, and a suitable solvent

-116-




with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1,  $R^c$  representing hydrogen or  $C_{1-4}$ alkyl,  $R^{c'}$  representing CN,  $NR^9R^{10}$ ,  $-C(=O)-NR^9R^{10}$ ,  $-C(=O)-C_{1-6}$ alkyl or  $R^7$ ;

- 5 n) reacting an intermediate of formula (XVIII) with  $CBr_4$  in the presence of a suitable catalytic salt, a suitable base and a suitable solvent




- o) reacting an intermediate of formula (XIV) with  $Cl_2C=S$  in the presence of a suitable solvent



with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;


- p) reacting an intermediate of formula (XV) with an intermediate of formula (XVI) in the presence of a suitable solvent

-117-



with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;



q) reacting an intermediate of formula (XXIX) with an intermediate of formula (XXX) in the presence of hydroxybenzotriazole and ethyldimethylaminopropyl carbodiimide, a suitable solvent and optionally in the presence of a suitable base



with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $R^9$ ,  $R^{10}$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1 and  $C_{2-6}alkenyl'$  representing  $C_{2-6}alkenyl$  optionally substituted with cyano;


r) reacting an intermediate of formula (XXXI) with an intermediate of formula (XXXII-1) or (XXXII-2) in the presence of hydroxybenzotriazole, ethyldimethylaminopropyl carbodiimide and a suitable solvent, and optionally in the presence of a suitable base

-118-



with  $\text{R}^1, \text{R}^2, \text{R}^4, \text{R}^{13}, \text{R}^{14}, \text{X}_1, m, n, -a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;


s) reacting an intermediate of formula (XI-b) with an intermediate of formula (XXXIII) in the presence of a suitable solvent



5

with  $\text{R}^1, \text{R}^2, \text{R}^4, \text{R}^{16}, \text{X}_1, m, n, -a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;

t) reductive methylation of an intermediate of formula (XXXIV) with formaldehyde in the presence of a suitable catalyst, a suitable reductive agent and a suitable solvent




10 with  $\text{R}^1, \text{R}^2, \text{R}^4, \text{X}_1, m, n, -a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;




-119-

u) reacting an intermediate of formula (XXXIV) with 2,5-dimethoxytetrahydrofuran in the presence of a suitable acid



with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;


- 5 v) reacting an intermediate of formula (XXXV) with an intermediate of formula (XXXVI) in the presence of  $n\text{BuLi}$  and a suitable solvent



with  $R^1$ ,  $R^2$ ,  $R^4$ ,  $R^7$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  as defined in claim 1;

- 10 and, if desired, converting compounds of formula (I) into each other following art-known transformations; and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or conversely, converting the acid addition salt form into the free base by treatment with alkali; and, if desired, preparing stereochemically isomeric forms,
- 15 *N*-oxide forms or quaternary amines thereof.
13. A product containing (a) a compound according to any one of claims 1 to 6, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in the treatment of HIV infection.
- 20 14. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound according to any one of claims 1 to 6, and (b) another antiretroviral compound.
- 25 15. A compound of formula

-120-




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined in claim 1 and  $W_3$

5 represents a suitable leaving group.


16. A compound of formula



a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined in claim 1.

17. A compound of formula




a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined in claim 1 and

$C_{2-6}alkenyl'$  represents  $C_{2-6}alkenyl$  optionally substituted with cyano.

18. A compound of formula




-121-

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein

$R^1$ ,  $R^2$ ,  $R^4$ ,  $X_1$ ,  $m$ ,  $n$ ,  $-a^1=a^2-a^3=a^4-$  and  $-b^1=b^2-b^3=b^4-$  are as defined in claim 1 and  $C_{2-6}alkenyl'$  represents  $C_{2-6}alkenyl$  optionally substituted with cyano.

5

19. A compound of formula



(III-b)

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine and a stereochemically isomeric form thereof, wherein  $R^4$  and  $X_1$  are as defined in claim 1.

10

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/08953

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7    C07D403/12    C07D401/12    C07D407/12    C07D409/12    C07D413/12 C07D417/12    C07D403/14    C07D233/96    A61K31/505    C07D239/46 C07D239/48    C07C211/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7    C07D    A61K    C07C		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) CHEM ABS Data, WPI Data, EP0-Internal		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 958 935 A (DAVIS JEREMY MARTIN ET AL) 28 September 1999 (1999-09-28) claim 1; examples 97,99 ---	1
X	EP 0 945 443 A (JANSSEN PHARMACEUTICA N.V., BELG.) 29 September 1999 (1999-09-29) cited in the application page 1, line 1 -page 2, line 14; claim 1 --- <div style="text-align: center;">-/--</div>	1-18
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  <div style="text-align: center;">18 November 2002</div>		Date of mailing of the international search report  <div style="text-align: center;">26/11/2002</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  <div style="text-align: center;">Samsam Bakhtiary, M</div>

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08953

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	D'AURIA M ET AL: "Photochemical Dimerization in Solution of Arylacrylonitrile Derivatives" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 53, no. 51, 22 December 1997 (1997-12-22), pages 17307-17316, XP004106551 ISSN: 0040-4020 page 17311 compound n° 7	19
Y	EP 1 002 795 A (JANSSEN PHARMACEUTICA NV) 24 May 2000 (2000-05-24) page 1, line 1 -page 2, line 9; claim 1	1-18
Y	WO 00 27825 A (AKEN KOEN JEANNE ALFONS VAN ;KOYMANS LUCIEN MARIA HENRICUS (BE); H) 18 May 2000 (2000-05-18) cited in the application page 1, line 1 -page 3, line 24; claim 1	1-18

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08953

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5958935	A	28-09-1999	AU 7631496 A EP 0862560 A1 WO 9719065 A1 US 6235746 B1	11-06-1997 09-09-1998 29-05-1997 22-05-2001
EP 0945443	A	29-09-1999	AU 751573 B2 AU 3599699 A BG 104738 A BR 9909191 A CA 2324919 A1 CN 1295564 T EE 200000532 A WO 9950250 A1 EP 1245567 A1 EP 0945443 A1 HR 20000620 A1 HU 0101204 A2 JP 2002509920 T NO 20004810 A PL 343196 A1 SK 14062000 A3 TR 200002760 T2 US 6197779 B1 US 2001011094 A1 EP 0945442 A1	22-08-2002 18-10-1999 30-04-2001 05-12-2000 07-10-1999 16-05-2001 15-02-2002 07-10-1999 02-10-2002 29-09-1999 30-06-2001 28-10-2001 02-04-2002 26-09-2000 30-07-2001 11-06-2001 21-12-2000 06-03-2001 02-08-2001 29-09-1999
EP 1002795	A	24-05-2000	EP 1002795 A1 AU 6200899 A BG 105418 A BR 9915552 A EE 200100252 A HR 20010161 A1 HU 0104177 A2 JP 2002529456 T NO 20011696 A PL 347586 A1 SK 6032001 A3 CN 1322198 T CZ 20011533 A3 WO 0027825 A1 TR 200101306 T2	24-05-2000 29-05-2000 30-11-2001 14-08-2001 15-10-2002 28-02-2002 28-03-2002 10-09-2002 04-04-2001 08-04-2002 07-01-2002 14-11-2001 17-10-2001 18-05-2000 22-10-2001
WO 0027825	A	18-05-2000	AU 6200899 A BG 105418 A BR 9915552 A CN 1322198 T CZ 20011533 A3 EE 200100252 A WO 0027825 A1 EP 1002795 A1 HR 20010161 A1 HU 0104177 A2 JP 2002529456 T NO 20011696 A PL 347586 A1 SK 6032001 A3 TR 200101306 T2	29-05-2000 30-11-2001 14-08-2001 14-11-2001 17-10-2001 15-10-2002 18-05-2000 24-05-2000 28-02-2002 28-03-2002 10-09-2002 04-04-2001 08-04-2002 07-01-2002 22-10-2001